



**UMY** UNIVERSITAS  
MUHAMMADIYAH  
YOGYAKARTA  
Unggul & Islami



4<sup>th</sup> International Conference of Sustainable Agriculture

# CERTIFICATE OF PARTICIPATION

This certificate is awarded to:

**Parwi**

in acknowledgement of your participation as

**PRESENTER**

at 4<sup>th</sup> International Conference of Sustainable Agriculture (ICoSA)  
held from 25<sup>th</sup> - 26<sup>th</sup> August 2021 at the Universitas Muhammadiyah  
Yogyakarta, Indonesia by virtual conference

Chairperson of  
the 5<sup>th</sup> ICoSI 2021




Dr. Yeni Rosilawati, SIP, SE., M.M

Chairperson of  
4<sup>th</sup> ICoSA 2021



Dr. Ihsan Nurkomar, S.P

Rector of Universitas  
Muhammadiyah Yogyakarta



Dr. Ir. Gunawan Budiyanto, M.P, IPM

ISSN 1751-1307

# **IOP** Conference Series

**Earth and Environmental Science**

1st Workshop on Metrology for Agriculture  
and Forestry (METRO-AGRIFOR)

# 275

VOLUME 275 – 2019

3–2 October 2018  
Asolo, Italy

**EDITOR**  
Enrico Pitta Tomassini

The open access journal for conference proceedings

[iopscience.org/jpcs](http://iopscience.org/jpcs)

**IOP** Publishing

PAPER • OPEN ACCESS

## Preface

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **985** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Preface](#)
- [PREFACE](#)
- [Preface](#)

### ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.  
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

**Learn more. Apply today!**

## PREFACE

The 4<sup>th</sup> ICoSA (International Conference on Sustainable Agriculture) 2021 is annual international scientific meeting organized by Department of Agrotechnology, Faculty of Agriculture, Universitas Muhammadiyah Yogyakarta (UMY). This conference is dedicated to facilitate multidisciplinary discussion among stakeholders (both domestic and/or international) related to the major issues found in agricultural sectors, particularly in the implementation of sustainable agriculture. This conference also mediates a beneficial exchange of information to support the improvement and the development of agriculture. For its fourth implementation, the 4<sup>th</sup> ICoSA 2021 brought the theme of “**Trend in Food Self-Sufficiency during Pandemic of COVID-19**”. Due to the global travel restrictions applied during the pandemic of COVID-19, this conference was conducted in the form of virtual meeting on August 25-26<sup>th</sup> 2021. Despite its virtual form of conference, we recorded high enthusiasm from the participants coming from different areas and institutions across Indonesia that applied to join in this conference. This excitement was reflected by 93 participants from Indonesia, Malaysia, Thailand, and the United States. Regarding its implementation, this conference provided two main activities, namely plenary talks and parallel discussion regarding various fields of studies related to agriculture.

After performing highly selective scientific review on the submitted paper, we are delighted to present this proceeding as a paper compilation representing various field of studies, such as crop production, crop protection, management of land use, and postharvest. We would like to express our gratitude to the organizing committees and supporting parties for their hard work and precious support, either during the conference and the post- conference publication, thus leading to a success of 4<sup>th</sup> ICoSA 2021.

We sincerely hope that this conference can be considered as a scientific forum providing high quality discussion among researchers and other related stakeholders in agricultural sectors. We belief that the proceeding may serve as useful source of reference for further studies which will contribute significantly for the future development of agricultural sectors.

The overall conference was organized by our organizing committees. On the other hands, the processing of this post-conference publication was be managed by a scientific committee, consisting of the editorial board and peer reviewers selected from various institutions from different countries and expertise. Editorial board plays an important role in setting the international standing and reviewing the quality of the submitted papers in order to fulfill the standard requested by the publishers. Peer reviewers representing each field of study play a critical role in reviewing the eligibility of the submitted papers based on its content.



**Organizing Committee of 4<sup>th</sup> ICoSA 2021****Steering Committee**

Dr. Gunawan Budiyo  
 Dr. Indira Prabasari  
 Dr. Innaka Ageng Rineksane  
 Dr. Lis Noer Aini  
 Dina Wahyu Trisnawati, Ph.D.

**Conference Chairman**

Dr. Ihsan Nurkomar

**Organizing Committee**

**Secretary** : Dr. Siti Nur Aisyah  
 Pradika Bunga, S.P.  
 Wafa Nadia R., S.P.

**Conference Program** : Taufiq Hidayat, S.P., M.Sc.

**Equipment** : Bambang Heri Isnawan, M.P.

**IT Support** : Takaful Pambudi Raharjo, S.P.  
 Husama Allauddin Bariq, S.P.

**Post-Conference Publication** : Dr. Siti Nur Aisyah

**Scientific Committee of 4<sup>th</sup> ICoSA 2021****Editorial Board**

**Prof. Damayanti Buchori**  
 IPB University, Indonesia  
 Expertise: Entomology  
**Scopus ID: 18041564100**

**Prof. Maria Rosa Mosquera-Losada**  
 University of Santiago de Compostela, Spain  
 Expertise: Agroecosystem and Soil Sciences  
**Scopus ID: 6506769402**

**Prof. Keith F. A. Walters**  
 Harper Adams University, United Kingdom  
 Expertise: Invertebrate Biology and Pest  
 Management  
**Scopus ID: 7102916144**

**Prof. Normah Mohd Noor**  
 Universiti Kebangsaan Malaysia, Malaysia  
 Expertise: Biotechnology and Tissue Culture  
**Scopus ID: 6602652954**

**Peer Reviewers**

**Dr. Lis Noer Aini**  
 Universitas Muhammadiyah Yogyakarta,  
 Indonesia  
 Expertise: Environment and Soil Sciences

**Dina Wahyu Trisnawati, Ph.D.**  
 Universitas Muhammadiyah Yogyakarta,  
 Indonesia  
 Expertise: Agroecology and Pest

---

**Scopus ID: 57204425510**

Management

**Scopus ID: 56503879400**

**Dr. Laily Ilman Widuri**

Universitas Jember, Indonesia

Expertise: Plant Physiology

**Scopus ID: 57196088116**

**Dr. Siti Nur Aisyah**

Universitas Muhammadiyah Yogyakarta,

Indonesia

Expertise: Plant Biotechnology

**Scopus ID: 57190760647**

**Dr. Ihsan Nurkomar**

Universitas Muhammadiyah Yogyakarta,

Indonesia

Expertise: Entomology

**Scopus ID: 57195399415**

**Dr. Eko Hanudin**

Gadjah Mada University, Indonesia

Expertise: Environment and Soil Sciences

**Scopus ID: 57196062767**

**Dr. Elly Syafriani**

Universitas Gadjah Mada, Indonesia

Expertise: Agronomy

**Scopus ID: 57189593893**

**Dr. Wakhid**

SMART Research Institute, Indonesia

Expertise: Entomology

**Scopus ID: 57196062767**

**Dr. Listihani**

Universitas Mahasaraswati Denpasar,

Indonesia

Expertise: Plant Virology

**Scopus ID: 57204636685**

---

PAPER • OPEN ACCESS

## Peer Review Statement

To cite this article: 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **985** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Peer review declaration](#)
- [Peer review declaration](#)
- [Peer Review Declaration Form Guidance](#)

### ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.  
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

**Learn more. Apply today!**

## Peer Review Statement

All papers published in this volume have been reviewed through processes administered by the Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

- **Type of peer review:** Double Anonymous
- **Conference submission management system:** Morressier
- **Number of submissions received:** 93
- **Number of submissions sent for review:** 75
- **Number of submissions accepted:** 59
- **Acceptance Rate (Submissions Accepted / Submissions Received × 100):** 63.4
- **Average number of reviews per paper:** 1
- **Total number of reviewers involved:** 13
- **Contact person for queries:**  
**Name:** Dr. Siti Nur Aisyah  
**Email:** sitinur@umy.ac.id  
**Affiliation:** Dept. of Agrotechnology, Faculty of Agriculture, Universitas Muhammadiyah Yogyakarta, Bantul 55183, Indonesia





- 
- OPEN ACCESS** 012029  
**Carrying Capacity for Land Rehabilitation Strategy in Merawu Watershed, Banjarnegara District**  
P Marcella, H Suryatmojo and L R W Faida  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012030  
**Application of Goat Manure Briquettes on Red Chili Cultivation in Coastal Sandy Soil**  
G Budiyanto, Mulyono and F Achmad  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012031  
**The Effect of The Application of Soil Biostructures Created Using Biochar and Seaweed Extract on Upland Rice Growth**  
L M H Kilowasid, S Samiri, M J Arma, L Sabaruddin, A Bahrin, L Afa, T C Rakian and W S A Hisein  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012032  
**Soil Chemical Properties and Microbial Biomass Respond during Land Use Change**  
**Parwi**, Muhammad, F D Dewanti and R Priyadashini  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012033  
**Effect of Different Soil Fertility on Growth and Development of Two Citrus Cultivars Under Two Locations**  
N F Devy, M E Dwiastuti and Hardiyanto  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012034  
**Land Suitability Re-evaluation for Fruit Crops in the Post-Eruption Merapi 2010**  
L N Aini, B H Isnawan and E Hanudin  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012035  
**Identification of Land Cover Types and Their Changes in Kutalimbaru Sub-district, Deli Serdang District, North Sumatra, Indonesia**  
Rahmawaty, H Kurniawan, M M Harahap and A Rauf  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012036  
**Location Determination of Critical Land Indicative as Basis for Forest and Land Rehabilitation in Merawu Watershed, Banjarnegara District**  
F B Ginting, H Suryatmojo and Widiyatno  
[+ Open abstract](#) [View article](#) [PDF](#)

PAPER • OPEN ACCESS

## Soil Chemical Properties and Microbial Biomass Respond during Land Use Change

To cite this article: Parwi *et al* 2022 *IOP Conf. Ser.: Earth Environ. Sci.* **985** 012032

View the [article online](#) for updates and enhancements.

You may also like

- [The role of soybean agroforestry in mitigating climate change in Indonesia](#)  
Djoko Purnomo, Maria Theresia Sri Budiastuti and Desy Setyaningrum
- [Compatibility of variety and planting pattern of Maize \(\*Zea mays\*\) in early Ganitri \(\*Elaeocarpus ganitrus\* Roxb.\) based agroforestry in dry land](#)  
A Rohandi and D Swestiani
- [Growth performance of sengon \(\*Falcataria mollucana\*\) and manglid \(\*Magnolia champaca\*\) at different spacing distance](#)  
A Hani and D Swestiani



**ECS** The Electrochemical Society  
Advancing solid state & electrochemical science & technology

### 242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Early hotel & registration pricing ends September 12

Presenting more than 2,400 technical abstracts in 50 symposia

The meeting for industry & researchers in  
**BATTERIES**  
**ENERGY TECHNOLOGY**  
**SENSORS AND MORE!**

 Register now!

 **ECS Plenary Lecture featuring M. Stanley Whittingham,**  
Binghamton University  
Nobel Laureate –  
2019 Nobel Prize in Chemistry



# Soil Chemical Properties and Microbial Biomass Respond during Land Use Change

Parwi<sup>1\*</sup>, Muhammad<sup>1</sup>, F D Dewanti<sup>2</sup>, R Priyadashini<sup>2</sup>

<sup>1</sup>Department of Agrotechnology, Faculty of Sciences and Technology, University of Darussalam Gontor, Indonesia

<sup>2</sup>Department of Agrotechnology, Faculty of Agriculture, Universitas Pembangunan “Veteran” East Java, Indonesia

\*E-mail: parwi@unida.gontor.ac.id

**Abstract.** In Ponorogo, East Java, Indonesia, pine-based agroforestry was mostly converted to sengon-based agroforestry before partly converted to monoculture. Land-use changes during the process affect soil properties. There is few information about the effects of land-use changes from pine-based agroforestry systems to sengon-based agroforestry systems and monocultures on soil chemical properties and microbial biomass. The aims of this study were to analyze the response of soil chemical properties and microbial biomass to land use changes in latosol Indonesian. Samples from three land-use changes, i. e. pine-based agroforestry, sengon-based agroforestry and monoculture were analyzed. Soil samples were taken at 2 depths, i. e. 0-30 cm and 30-60 cm. As the results, it was found that the land-use change affected the chemical properties and microbial biomass of the soil. Land-used change from pine-based agroforestry to sengon-based agroforestry have no significant effect on pH, organic carbon, available P, and microbial biomass in 0-30 cm depth. Meanwhile, in 30-60 cm depth, microbial biomass in sengon-based agroforestry was higher than that in pine-based agroforestry. The pine-based agroforestry which was converted into monoculture, reduced the soil organic carbon and available P concentration, contrast with the soil microbial biomass which was increased in 0-30 cm depth. The highest soil microbial biomass was found in sengon-based agroforestry in 30-60 cm depth. The conclusion of this study implicated that land use in monoculture decreased soil fertility. Therefore, conversion of pine-based agroforestry to monoculture is full of risks.

**Key words:** Soil chemical, microbial, land-use

## 1. Introduction

Pine agroforestry systems have been developed in East Java, especially areas that have an altitude above 800 m above sea level. Pine plants can contribute organic carbon to the soil through leaves and roots. The amount of organic carbon donated to the soil depends on the age of the pine, the older the pine, the greater the organic carbon given to the soil [1,2]. Pine plants can affect the chemical and biological properties of the soil [3]. Pine plants besides being able to be planted in monoculture, they can also be planted in agroforestry. The problem with pine-based agroforestry was that it takes too long to harvest pine trees, which was more than 10 years. An alternative to pine plants was sengon, with the advantage of relatively shorter harvest times compared to pine.

Sengon is a legume that grows very fast. This was related to the ability of sengon which can be in symbiosis with rhizobium. Rhizobium in symbiosis with rhizobium plays a role in remembering N from



the air [4]. Rhizobium bacteria can donate nitrogen to sengon plants. The very fast growth of sengon was beneficial in donating organic carbon to the soil. Sengon plant leaves that fall to the ground and rotting roots are a source of organic carbon in the soil [5].

Pine-based agroforestry land in addition to being converted into sengon-based agroforestry land, there was also conversion to monoculture land. Monoculture lands have different management management from agroforestry systems. In monoculture land, the input of organic matter depends on the amount of organic fertilizer applied to the soil. The quality of organic fertilizer depends on the source of organic matter and its quality. The quality of organic fertilizers can be determined based on the C/N ratio [6]. The quality of organic fertilizer can determine the addition of organic carbon into the soil.

Changes in organic carbon from pine-based agroforestry systems to sengon agroforestry and monocultures have the potential to affect soil chemical properties and microbial biomass. Microbial biomass in agroforestry systems was influenced by the type of annual plant. Microbial biomass in Acacia-based agroforestry > Populus-based agroforestry > Eucalyptus-based agroforestry in alkaline soils of India [7]. The availability of scientific articles that discuss changes in organic carbon due to land conversion was very limited. This study aims to examine changes in soil chemical properties and microbial biomass due to changes in land use. This information was useful for assessing the impact of land use change

## 2. Materials and Methods

### 2.1. Sampling

The research was conducted in Mrayan Village, Ngrayun District, Ponorogo Regency, East Java, Indonesia. Samples in the form of soil will be taken randomly from the rhizosphere of porang plants in agroforestry systems (pine and sengon) and monoculture. Each sampling site was repeated randomly for 3 replications. For each replication, 5 sub samples were taken randomly, so that the number of sub samples taken was  $3 \times 3 \times 5 = 45$  sub samples. Soil samples were taken at a depth of 0-30 cm and 30-60 cm. Sub samples for each planting and the same replication were composited. The composite results were taken 2 kg of soil and divided into 2 parts, each 1 kg. One part was put in a paper bag and stored at room temperature prior to soil chemical analysis. Another part was put into a paper bag and stored in the refrigerator for analysis of soil microbial biomass. Soil analysis and microbial biomass were carried out at the Agrotechnology Laboratory, university of Darussalam Gontor.

### 2.2. Soil chemical analysis.

Soil chemical analysis will be carried out on ground samples with a size of < 2 mm. Organic C analysis was carried out by the Black and Walkey method [8]. Available P analysis was using the Olsen method [9]. Soil pH was determined with water solvent (1:5).

### 2.3. Microbial biomass analysis.

Microbial biomass analysis was carried out using the fumigation method [10]. Microbial biomass was determined by calculating the difference in organic carbon content between after fumigation and before fumigation, then the result was converted into microbial biomass by multiplying by 0.35.

### 2.4. Statistical analysis

Data analysis will be carried out using one-way analysis of variance (ANOVA) using the SPSS program. LSD (Student's t lest significant difference) analysis was used to compare between treatments.

## 3. Results and Discussions

### 3.1. Soil chemical properties

At a depth of 0-30 cm, it was found that sengon-based agroforestry had no different organic pH and organic carbon compared to pine-based agroforestry and monocultures. The average soil pH was 6.28-6.81, while the average soil organic C is 1.29-1.49%. AvailableP in agroforestry systems were higher

than in monocultures (Table 1). In pine-based agroforestry and sengon can contribute organic carbon into the soil through leaf fall in the top layer of soil. Leaves that have touched the ground will decompose to produce organic P and organic acids. Organic acids were able to release P which is bound by the soil so that the available P content increases. This situation supports the research of Wang et al. [6] which states that agroforestry lands have higher available P levels compared to monocultures.

**Table 1.** Soil chemical properties

	Pine-based agroforestry	Sengon-based agroforestry	Monoculture
Depth: 0-30 cm			
pH	6,28±0,56 a	6,81±0,34 a	6,35±0,04 a
Organic Carbon (%)	1,41±0,33 a	1,49±0,09 a	1,29±0,09 a
Available P (ppm)	24,43±1,23 b	28,11±4,61 b	13,64±1,72 a
Depth: 30-60 cm			
pH	5,19±0,83 a	6,31±0,39 b	5,31±0,47 a
Organic Carbon (%)	0,54±0,19 b	0,56±0,07 b	0,49±0,20 a
Available P (ppm)	24,35±0,62 b	23,89±0,38 b	13,50±0,87 a

Data followed by the same letter in the same column were not significantly different based on the 5% LSD test

At a depth of 30-60 cm, it was found that soil pH in sengon-based agroforestry was higher (21.57%) than pine-based agroforestry (5.19). Sengon was a legume plant that contains many nutrients including calcium which can increase soil pH. Organic matter that has a C/N of 27 can increase soil pH higher than organic matter that has a C/N of 15 [6]. The monoculture system had a lower pH (15.85%) than sengon-based agroforestry system (6.31). In monoculture land, the addition of organic matter was relatively small compared to agroforestry systems. Decomposition of organic matter will increase soil pH. Organic matter can increase pH in Typic Paleudalf soil [11].

There was not significant difference between organic carbon levels in agroforestry systems. The average organic carbon in agroforestry systems is 0.54-0.56%. This was related to the contribution of plant roots to soil organic carbon. Roots that have rotted will provide organic carbon input into the soil. In addition, with active plant roots, soil microbes will develop better. Dead microbes were a source of organic carbon in the soil. The monoculture system had lower organic C (9.26% and 12.50%) than the pine-based agroforestry (0.54%) and sengon-based agroforestry (0.56%). In monoculture porang plants, plant roots will focus at a depth of 0-30 cm, a few reach a depth of 30-60 cm. Coffee agroforestry systems have a much higher SOC than monocrop coffee [12]. Likewise, Hergoual'h et al. [13] found that coffee agroforestry systems with leguminous tree species had higher soil organic carbon than monoculture coffee.

Pine-based agroforestry land had higher available P (44.91%) compared to porang land in monoculture (13.50 ppm). The highest available P levels were in the Pine-based agroforestry area and the lowest was in the monoculture system (Table 1). The pine plant has the characteristics of leaves that contain a lot of lignin so that pine leaves were difficult to decompose. In organic matter that was difficult to decompose, the decomposition product is in the form of low nutrients but produces organic acids that were able to release P which was absorbed by the soil so that it becomes available P in the soil. The increase in total soil organic carbon due to the addition of organic matter can increase the available P in Xerofluent [14]. This fact was different from Emiru and Gebrekidan [15] which stated that the decrease in soil organic carbon was not followed by a decrease in available P in soil in Senbat Watershed, Western Ethiopia.


### 3.2. Microbial biomass

At a depth of 0-30 cm, microbial biomass in sengon-based agroforestry was higher than pine-based agroforestry (Figure 1). This is related to the source of energy and nutrients needed by soil microbes. Pine leaves have different characteristics from sengon leaves. Sengon leaves contain better nutrients

than pine leaves so that the development of microbes is higher. Microbial biomass in agroforestry systems depends on the type of annual crop. *Ficus hookerii*-based agroforestry (801 mg kg) was higher than *Terminalia myriocarpa*-based agroforestry (790 mg kg) [16].

Monoculture system had higher microbial biomass than sengon-based agroforestry and pine-based agroforestry (Figure 1). This result is different from the results of research by Lepcha et al. [17], that Cardamom-based agroforestry has higher microbial biomass than rice plants in monoculture. This occurs due to differences in the input of organic matter into the soil. Microbial biomass is related to the concentration of nutrients in the soil. Microbial biomass is more determined by the concentration of N than P [18]

At a depth of 30-60 cm, sengon-based agroforestry systems were not different from pine-based agroforestry. This means that the root system of the two types of plants has not effect on soil microbial biomass. Monoculture systems have lower microbial biomass than sengon-based agroforestry. Porang plants grown in monoculture have shallow roots so that the contribution of organic matter from the roots was low at a depth of 30-60 cm.



**Figure 1.** Microbial biomass

### 3.3. Correlation soil chemical properties and microbial biomass

At a depth of 0-30 cm, microbial biomass was negatively correlated with available soil P (Table 2). Microbial biomass increases with decreasing soil available P. This result was in line with the research by Fujita et al. [18] which revealed that microbial biomass was negatively correlated with soil available P. However, these results differ from those of Lepcha et al [17], in that microbial biomass was positively correlated with soil P available in the Eastern Himalayas. Furthermore, Tan et al. [19] stated that microbial biomass was positively correlated with available P in Boreal Aspen Forest soil.

**Table 2.** Correlation soil chemical properties and microbial biomass

	pH	Microbial biomass	Organic Carbon	Available P
Depth 0-30 cm				
pH	1	-.052	.837**	.466
Microbial biomass	-.052	1	-.306	-.776*
Organic Carbon	.837**	-.306	1	.465
Available P	.466	-.776*	.465	1
Depth 30-60 cm				
pH	1	.613	.001	.333
Microbial biomass	.613	1	.287	.644
Organic Carbon	.001	.287	1	.780*
Available P	.333	.644	.780*	1

Soil pH was positively correlated with soil organic carbon (Table 2). Soil pH increases with increasing soil organic carbon. This was in line with Kaiser et al. [20] that soil pH was positively correlated with soil organic carbon. Soil pH was positively correlated with soil organic carbon in European agricultural soils [21].

Meanwhile, at a depth of 30-60 cm, soil organic carbon was positively correlated with available P (Table 2). Increased organic carbon levels will cause available P to increase as well. Organic carbon was positively correlated with soil available P in Ultisol soils in Owerri, Southeastern [22].

#### 4. Conclusions

Land use change affected the chemical properties and microbial biomass of the soil. Changes from agroforestry to monoculture decreased available P, in depth 0-30 cm and decreased available P and organic C in depth 30-60 cm. Changes from pine-based agroforestry to monoculture caused microbial biomass increase in depth 0-30 cm. Pine-based agroforestry converted to sengon-based agroforestry can be increased pH in depth 30-60 cm and microbial biomass in depth 0-30 cm. Results of this study implicated that monoculture agricultural land use was worse than sengon-based agroforestry.

#### Acknowledgments

We thank the University of Darussalam Gontor for funding this research in 2021.

#### References

- [1] Cheng X, Han H, Kang F, Song Y & Liu K 2013 *Journal of Plant Interactions* **9** 521–528.
- [2] Helmisaari HS, Makkonen K, Valtonen E, Kellomaki S, Malkonen F 2002 *Forest Ecology and Management* **165** 317–326.
- [3] Mosculo A, Sidari M, Mercurio R 2007 *Forest Ecology and Management* **242** 412–418.
- [4] Wedhastri S, Prahastiwi Y, Widada J, Widiyanto D and Kabirun S 2013 *Journal of Biotechnology* **18** 64–69.
- [5] Agus C, Putra PB, Faridah E, Wulandari D & Napitupulu RRP 2016 *Procedia Engineering*, 159329–337.
- [6] Wang L, Li Y, Prashe SO, Yan B, Ou Y, Cui H & Cui Y 2019 *Bioresource Technology* **289** 121745.
- [7] Kaur B, Gupta SR & Singh G 2000 *Applied Soil Ecology* **15** 283–294.
- [8] Walkey A and Black IA 1934 *Soil Sci.* 3739–28.
- [9] Olsen SR, Cole CV, Watanabe FS and Dean LA 1954 Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Dept Agric Circ, pp. 939.
- [10] Vance ED, Brookes PC and Jenkinson DS 1987 *Soil Biology and Biochemistry* **19** 703 – 707.
- [11] Pietri AJC & Brookes PC 2008 *Soil Biology and Biochemistry* **40** 1856–1861.
- [12] Tumwebaze SB, Byakagaba P 2016 *Agriculture, Ecosystems and Environment* **216** 188–193
- [13] Hergouale'h K, Blanchart E, Skiba U, Hénault C, Harmand JM 2012 *Agric. Ecosyst. Environ.* **148** 102–110.
- [14] Melero S, Porras JCR, Herencia JF & Madejon E 2006 *Soil and Tillage Research* **90** 162–170.
- [15] Emiru N and Gebrekidan H 2013 *ARPJN Journal of Agricultural and Biological Science* **8** 206–212.
- [16] Singh M, Das SK and Avasthe RK *Indian J. of Agroforestry* **20** 25–29.
- [17] Lepcha NT, Devi NB 2020 *Ecol Process* **9** 10–14.
- [18] Fujita K, Miyabara Y & Kunito T 2019 *European Journal of Soil Biology* **91** 1–8.
- [19] Tan X, Chang S & Kabbzem R 2008 *BiolFertil Soil* **44** 471–479.
- [20] Kaiser EA, Mueller T, Joergensen RG, Insam H & Heinemeyer *Soil Biology and Biochemistry* **24** 675–683.
- [21] Xu H & Zhang C 2020 *Science of The Total Environment* **14** 1–11.
- [22] Okon MA, Osuji GE, Uzoho BU, Ahukaemere CM 2014 *International Journal of Research in Agriculture and Forestry* **1** 18–22.