



CERTIFICATE

Nomor : DL.02.01/1/10488/2021

This Certificate Is Proudly Presented To

Rindang Diannita
as
PRESENTER

The 1st International Conference on Nursing and Public Health (ICoNPH) 2021
which is part of the 4th International Conference of Health Polytechnic Surabaya (ICoHPS) 2021
“Empowering community and health professionals in mitigating the impacts of covid-19 pandemic”
Surabaya, 6-7 October 2021

Director of
Poltekkes Kemenkes Surabaya



drg. Bambang Hadi Sugito, M.Kes

General Chair

Dr. Siti Nur Kholifah, SKM, M.Kep.Sp.Kom

Co Host :





Proceeding ICoHPS 2021

International Conference on Health Polytechnic Ministry of Health Surabaya

Poltekkes Kemenkes Surabaya

Health Polytechnic of the Ministry of Health Surabaya - Indonesia

Surabaya, 6 - 7 October 2021



ISBN 978-623-97447-4-8



9 786239 744748

**LIST OF INTERNATIONAL CONFERENCE COMMITTEE
HEALTH POLYTECHNIC OF SURABAYA IN 2021**

Director : drg. Bambang Hadi Sugito, M.Kes.

Person in charge :

1. Dr. Khambali, S.T., MPPM (Deputy Director I)
2. Dr. Hilmi Yumni, S.Kep., Ns., M.Kep., Sp. Mat (Deputy Director II)
3. Moch. Najib, S.Kp, M.Sc. (Deputy Director III)
4. Setiawan, S.KM., M.Psi (Head of the Center for Research and Community Service)

Steering Committee :

1. Dr. Sri Utami, SKp., M.Kes. (Head of the Center for Educational Development)
2. Dr. I Dewa Gede Hari Wisana, ST., MT. (Head of Quality Assurance Center)

Chairman : Dr. Siti Nur Kholifah, SKM, M.Kep.Sp.Kom

Vice Chairman 1 : Dr. Yessy Dessy Arna, M.Kep.Sp.Kom

Vice Chairman 2 : Dr. Dra. Anik Handayati, M.Kes

Secretary : 1. Sari Luthfiah, S.Kep.Ns, M.Kes
2. Heri Sumasto, S.Kep., M.M.Kes

Treasurer : 1. Fitri Rokhmalia, SST, M.KL
2. Tri Koerniasih, SE

A. Chair of Technical Program : Dr.Ir. Juliana Christyaningsih, M.Kes

1. Electro Medical Engineering Department

Coordinator : Dr. Triwiyanto, MSi

Member :

1. Dr. Endro Yulianto, MT
2. Andjar Pudji, ST.MT

2. Nursing Department

Coordinator : Dr. Padoli, SKp, M.Kes

Member :

1. Dr. Supriyanto, SKp, M.Kes
2. Dr. Dhiana Setyorini, M.Kep.Sp.Mat

3. Minarti, M.Kep.Sp.Kom
4. Adin Mu'afiro, SST, M.Kes
5. Hepta Anugraheni, S.Kep.Ns, M.Kep
6. Kusmini Suprihatin, SKp, M.Kep.Sp.A
7. Ach. Arfan Adinata, S,Kep,Ns., M.Kep

3. Midwifery Department

Coordinator : Evi Pratami, S.ST, M. Keb

Member :

1. Astuti Setiyani, SST., M.Kes
2. Rekawati Susilaningrum, M.Kes
3. Teta Puji Rahayu, SST., M.Keb
4. Deasy Irawati, SST., M.Keb
5. Ari Tri Rahayu, S.Keb

4. Medical Laboratory Technology Department

Coordinator : Ayu Puspitasari, S. T, M. Si

Member :

1. Wisnu Istanto, S. Pd M.PBI
2. Anita Dwi Anggraini, SST M. Si
3. Suhariyadi, S. Pd M. Kes
4. Pestariati, S. Pd, M. Kes
5. Retno Sasongkowati, S.Pd., S.Si., M. Kes

5. Environmental Health Department

Coordinator : Dr. Sri Poerwati, ST., M.Si

Member :

1. Winarko, SKM.,M.Kes
2. Ferry Kriswandana, S.ST.,MT
3. Irwan Sulistio, SKM.,M.Si
4. Hery Koesmantoro, ST.,MT

6. Nutrition Department

Coordinator : Inne Soesanti, S.SosS.Si., M.Kes

Member :

1. Taufiqurrahman, SKM, MPH
2. Ani Intiyati, SKM., M.Kes
3. Nur Hatijah, SKM., MKes
4. Erlyn Jayeng, SST., M.Gz

5. Nurul Hindaryani, S.Pd., M.Gz

7. Dental Nursing Department

Coordinator : Imam Sarwo Edi, S.Si.T, MPd

Member :

1. drg. Ida Chairanna Mahirawatie, M.Kes.
2. drg. Sri Hidayati, M.Kes.
3. Sunomo Hadi, S.Si.T., M.Kes.
4. Silvia Prasetyowati, S.Si.T., M.Kes.
5. Endang Purwaningsih, S,Si.T., SH., M.Pd.

B. Chair of Publication :Yohanes Kambaru Windi, MPH, PhD

1. Electro Medical Engineering Department

Coordinator : M.Ridho Makruf, ST., M.Si

Member :

1. Bedjo Utomo, M.Kes
2. Triana Rahmawati, ST.M.Eng
3. Dyah Titisari, ST., M.Eng.

2. Nursing Department

Coordinator : Dr. Anita Joeliantina, M.Kes

Member :

1. Dr. Jujuk Proboningsih, SKp, M.Kes
2. Dr. Moch. Bahruddin, M.Kep.Sp.KMB
3. Nikmatul Fadila, M.Kep.
4. Hasyim As'ari, S.Kep.Ns, M.Ked
5. Dyah Wijayanti, M.Kep.

3. Midwifery Department

Coordinator : Dr. Ns. Heru Santoso Wahito Nugroho, S.Kep, M.M.Kes

Member :

1. Suparji, SST., M.Pd., SKM
2. Dwi Purwanti, S.Kp., SST., M.Kes
3. Dwi Wahyu Wulan S, SST., M.Keb
4. Queen Khoirun Nisa Mairo, SST., M.Keb
5. Kharisma Kusumaningtiyas, S.SiT., M.Keb
6. Suryaningsih, S.SiT., M.Keb
7. Lilik Triyawati, A.Md.Keb., S.Pd., M.Kes

4. Medical Laboratory Technology Department

Coordinator : Evy Diah Woelansari, S.Si, M.Kes

Member :

1. Tacik Idayanti, S. ST
2. Ratno Tri Utomo, S. ST
3. Lully Hanni E, S. Si., M.Farm, Apt
4. Era Fitria Y., S.TrKes
5. Suliati, S. Pd., S.Si., M.Kes
6. Marlinda, A. Md. Kes

5. Environmental Department

Coordinator : Dr. Budi Yulianto, S.Pd, M.Kes

Member :

1. Dr. Ir. Iva Rustanti, E.W., M.T.
2. Marlik, S.Si., M.Si
3. Narwati, S.Si., M.Kes
4. Aries Prasetio, S.Si.,M.Si
5. Pratiwi Hermiyanti, S.ST.,M.KL

6. Nutrition Department

Coordinator : Annas Buanasita, SKM., M.Gz

Member :

1. Nuning Marina Pengge, SKM., M.Kes
2. Mujayanto, SKM., MPH
3. Melina Sari, M.Si
4. Riezky, S.ST., M.Gz

7. Dental Nursing Department

Coordinator : Isnanto, SSiT, M.Kes

Member :

1. drg. Soesilaningtyas, M.Kes.
2. Agus Marjianto, S.Si.T.,SKM.,M.Kes.
3. Siti Fitria Ulfah, S.Si.T.,M.Kes.
4. Mokhamad Ainu Riziq, S.ST.
5. Sanda Astrinata, S.ST.

C. Chair of Publicity : Syaifuddin, ST., MT

1. Electro Medical Engineering Department
2. Nursing Department

: Syevana, SST

: Eko Rustamadji, M.Tr.Kep

3. Midwifery Department : Titi Maharrani, M.Keb
4. Medical Laboratory Technology Department : Christ Kartika R., ST, M.S;
Riya Agustin, S.ST.
5. Environmental Health Department : Ernitasari, SST, MKL
6. Nutrition Department : Devi Eka Ratnasari, SST
7. Dental Nursing Department : Idot Suryopratomo, SE

D. Chair of Web and IT : Dr. Joko Suwito, SKp, M.Kes

1. Dr. Dwi Ananto Wibrata, SST, M.Kes
2. Li'ulliyah, S.Kom., M.Kom.
3. Putu Widiarsa Kurniawan, S.Kom
4. Aby Yazid Al Bustomi, M.Kep.Sp.KMB
5. Vicky Anjar Rizki Pratama, S.Kom

Secretariat

Coordinator : Yaimin, S.Pd, M.Pd, M.A.Ed

Member :

1. Edi Yulianto, S.M
2. Ira Rahayu Tyar Sari, S.ST

DAFTAR ISI

Overview Of Behavior And Control Of Dengue Fever In The Work Area Of The Sabokingking Palembang Health Center Year 2020	200-211
Farhani Sucitami, Ghina Ajeng Felicia, Emmy, Priyadi, Maksuk	
The Potential of Fermented Oyster Mushroom Filtrate (<i>Pleurotus ostreatus</i>) as Host-Seeking Attractant of <i>Aedes aegypti</i> Mosquito	212-220
Feby Carira Sindy, Pratiwi Hermiyanti, Fitri Rokhmalia, Demes Nurmayanti	
Management of Hazardous and Toxic (B3) Covid-19 Infectious Waste in Nganjuk Hospital 2021	221-226
Ahlun Najaa Nazzun Priyono Putro, Fitri Rokhmalia, Darjati	
Anxiety and Sleep Quality among Medical Workers in Harjono S Hospital	227-233
Dian AfifArifah, RatihAndhika Akbar R, Rindang Diannita	
The Effect Of Distance Of Benowo Waste Final Disposal (Tpa) Surabaya City On H2s And Nh3 Levels In The Settlement Environment	234-236
Imilda Lidiawati, Khambali, Darjati	
Relationship Of Hygiene And Sanitation With Microbiological Quality Of Drinking Water Depo Water During The Covid-19 Pandemic In 2021 (In Dupak Village, Krembangan District, Surabaya City)	237-238
M. Lazzuardhi Ilmi, Khambali, Demes Nurmayanti	
Influence Of So2 And Nh3 Gas Levels On Public Health Around Benowo Final Disposal Site, Surabaya City	239-244
Rifka Anggraeni, Khambali, Narwati	
The Effect Of Noise Due To Aircraft Passing In Settlements Around Juanda Airport On Community Subjective Complaints (Case Study in Sedati Gede Village, Sidoarjo in 2021)	245-249
Arina Khusnal Hidayaty, Khambali, Rachmaniyah, Demes Nurmayanti	
The Effect Of Working Period, Nutritional Status, And Smoking Habits On Work Fatigue At Pt. Atlantic Anugrah Metalindo Surabaya Year 2021	250-255
Presillia Aulina, Khambali, Suprijandani, Winarko, Rusmiati	
Society Role in Drinking Water Treatment with <i>Ipomoea carnea</i>	256-261
Sri Poerwati, Aries Prasetyo, Waella Septamari Budi	
<i>Ipomoea carnea</i> Leaf Extract As Antibacterial Drinking Water Deep Wells	262-266
Sri Poerwati, Aries Prasetyo, Waella Septamari Budi	
The Examination of The Quality of Chitosan from Bamboo Shell Waste with Variations of NaOH Concentration in the Deacetylation Proses	271-279
Sayyidah Nafysah Ahmad, Darjati, Hadi Suryono	
Analysis Of The Risk Of Dust/Particulate Matter (PM2,5) On Public Health (Case Studi : Place Of Manufacture Of Bricks In Kaloran Village, Ngronggot Sub-District, Nganjuk District On 2021)	280-287
Dzahabiyah Dwi Putri Ridayanti, Khambali, Hadi Suryono	
The Relationship Between Work Attitude With Work Fatigue at PT. “IK” Madiun Finishing Division 2021	288-293
Fheby Cahyati Nababan, Budi Yulianto, Sunaryo, Vincentius Supriyono	

Anxiety and Sleep Quality among Medical Workers in Harjono S Hospital

Dian Afif Arifah^{1(CA)}, Ratih Andhika Akbar R¹, Rindang Diannita¹

^{1(CA)}Department Occupational Safety and Health, Faculty of Health, Universitas Darussalam Gontor, Indonesia; dianafif@unida.gontor.ac.id (Corresponding Author)

ABSTRACT

Anxiety rate is widely increased and has become the most common symptom experienced by medical staff during the COVID-19 pandemic since the end of 2019. Recent data referred that more than 50% of medical staff experience moderate to high levels of anxiety. Sleep quality is variable that is highly associated with the human physical condition including anxiety. A person with good sleep quality is less likely to feel stressed, anxious, depressed, or experience physical fatigue. This study aims to determine the anxiety level among medical staff based on their sleep quality.

Forty-eight (48) medical staff were involved as respondents in this study which was taken randomly using a simple random sampling method. The independent variable is the sleep quality index (SQI), which is measured using the Pittsburg Sleep Quality Index (PSQI) instrument. While the dependent variable is the depression score which is measured using the Zung Self-rating Scale (SAS) instrument. Comparative analysis was used to compare anxiety scores within 2 groups based on SQI categories using an independent t-test test using 95% of Confidence Interval (CI).

Comparison test between 2 groups of sleep quality shows a significant difference (p-value/sig. 0.20) with a mean difference of 5,78. Based on the maximum score of anxiety, it can be concluded that HCW with good sleep quality has a 7.35 % lower level of anxiety compared to those who had worse sleep quality.

Keywords: *anxiety, sleep quality, health car workers.*

INTRODUCTION

The COVID-19 pandemic has widely increased workload for Health Care Workers (HCW) (1) which could causing impact on the quality of services and productivity (2). This situation will affecting not only the physical but also the mental health of HCWs (3). According to Labrague (2021) (4) anxiety is the most common symptom experienced by medical taff during the COVID-19 pandemic since the end of 2019. Recent data referred that more than 50% of medical taff experience moderate to high levels of anxiety. The rate were significantly higher than those observed in the general population (5–7).

While a low level of anxiety is beneficial to motivate, maintain self-defense and improved safety behavior, higher level of anxiety may have negative consequences on the mental-psychological health and work performance among workers. Several studies have found that the negative effects of a higher level of anxiety, including neural disorders, dizziness, sleep disturbance, vomiting and nausea (8,9). Extreme level of anxiety levels were also associated with impaired body function, negative coping mechanisms (such as increased intake of alcohol or drugs) stress, depression and increased burnout risk even suicidal ideation (10–12).

Factors related to the pandemic that are often associated with the physical problems of medical staff are intensive patient care, high patient mortality, high workload, role conflict and lack of time available to meet patient needs, etc. (4,13). Issues such as social stigmatization, shortage of personal protection equipment supplies, and heavy workload on the staff can aggravate this situation.

These condition will lead to serious consequences for both patients and the healthcare professionals (2). Several systematic reviews have found that high levels of burnout in health care professionals are associated with less-safe patient care. These consequences impose immense costs on the society (14,15).

Sleep quality is variable that is highly associated with human physical condition (16). It has been estimated that 90% of patients with depression complain about sleep quality (17). A person with good sleep quality is less likely to feel stressed, anxious, depressed or experience physical fatigue (18–20). For workers, having good quality of sleep is very important to maintain their physical and psychological condition while working. According to Arifah (21), the sleep quality index affects the level of fatigue experienced by medical taff who work in shifts. According to the study, workers with night shifts can maintain the quality of their sleep to prevent the risk of fatigue that occurs due to shift work. A workers with good sleep quality tends to feel

relaxed and stable so as to minimize the risk of psychological disorders such as stress, depression and anxiety.

Based on M Söderström(22), insufficient sleep can be a used as clinical indicator of burnout. Giorgi's(23)also proves that personal burnout was significantly associated with sleep disturbance among shift-work nurses.

Harjono Hospital is one of the Covid-19 care and isolation center in Ponorogo district. It provides isolation room with a capacity that can accommodate about 35 patients. However, the increasing number of cases caused the number of COVID-19 patients being overloaded. The Emergency Unit is a room for emergency patients entrance which operating 24 hours a day, 7 days a week. Patients received in emergency unit usually had acute cases occur suddenly or by accident without warning and require prompt and appropriate help. Due to the emergency unit operations require high accuracy that highly depends on human resources. High performance of medical staff in the emergency unit needed to provide health care services whatever patient's condition is. The risk of receiving patients with COVID-19 infection is high during pandemic.

This study aims to analyze anxiety experienced by Medical staff in the emergency Unit of RSUD dr. Harjono based on the their sleep quality.

METHODS

This study is an analytic observational study using a cross-sectional design. Data were collected in February to March 2021 at Dr. Hospital. Harjono S. Ponorogo. Fourty eight (48) medical staff involved as respondents in this study which taken randomly using simple random sampling method. Before the data collection, respondents were given an explanation and the research procedures and voluntarily filled out informed consent form. The collection procedure is described in the following chart:




Figure 1. Data collecting and analysis procedure

The independent variable is the sleep quality index (SQI)(24), which is categorized into ; "Good" (score <5) and "Bad" (score >5) based on the score of 7 indicators measured using the Pittsburg Sleep Quality Index (PSQI) instrument. While the dependent variable is the depression score which is measured using the Zung Self-rating Scale (SAS) instrument(25).

Data were both descriptive and analytically analysed. Microsoft Excel and Statistical Package for Social Sciences Software (SPSS) version 24.0 was used in all step of analyses. Comparative analysis will be used to compare anxiety scores within 2 groups based on SQI categories using independent t-test test using 95% of Confidence Interval (CI).

RESULTS

Table 1 described the Sleep experience in the last month and average depression score among 48 respondents. There are 7 components as indicators for sleep quality which indicates the symptoms experienced by average respondents in percentage. Overall sleep quality classified as "Good" or "Bad" based on the score in the 7 indicators.

Table 1. Respondent Characteristics based on Sleep quality and Anxiety Indicators

Indicators	Percentage
Sleep Quality Indicators	
Subjective Bad Sleep Quality	28,47 %
Sleep Latency*Frequence	29,16 %
Sleep Duration	34,02 %
Sleep Efficiency	29,16 %
Sleep Disturbance	17,36 %
Drugs Usage	0,00 %
Day Dysfunction	27,77 %
Anxiety Indicators	
Feel more nervous and anxious than usual	16,67%
Feel afraid for no reason at all	26,39%
Get upset easily or feel panicky	20,83%
Feel like falling apart	30,56%
Feel that something bad will happen	31,94%
Arms and legs shakedand trembled	19,44%
Bothered by headaches and back pain	29,17%
Feel weak and get tired easily	25,00%
Can't feel calm and hard to sit still	27,78%
Feel heart beating fast	19,44%
Bothered by dizzy spells	25,00%
Fainting spells or felt like it	9,72%
Can't breathe in and out easily	6,94%
Get numbness and tingling in the tingers and toes	15,28%
Bothered by stomach aches or indigestion	22,22%
Frequently urinate	19,44%
Hands are usually dry and warm	13,89%
Face gets hot and blushed	16,67%
Can't fall asleep and rest easily	25,00%
Had nightmares	18,06%

Table 1 describes the symptoms experienced by respondents based on indicators to assess sleep quality and anxiety levels on the PSQI and SAS instruments. Sleep problems that are most often experienced by medical personnel are lack of sleep duration (34.02%). A total of 54.17% of respondents with sleep problems stated that their sleep duration was less than 7 hours/day. Based on anxiety symptoms, most or 31.94% of medical personnel felt “Feel that something bad will happen”, “Feel like falling apart” (30.56 %) and “Bothered by headaches and back pain” (29.17 %). In general, the comparison of the Average anxiety score based on sleep quality is described in the table 2.


Table 1. Average Anxiety Score based on Sleep Quality

Sleep Quality	N	Percentage	Average Anxiety Score
Good	33	72,9 %	31.12
Bad	15	27,1 %	37.00

Σ	48	100 %	32.95
----------	----	-------	-------

Based on table 2, as many as 27.1% of medical personnel have bad sleep quality and an average anxiety score of 32.95. Medical personnel with bad sleep quality have an average anxiety score of 37.00 or 7.35% smaller than medical personnel with good sleep quality. Figure 1 described the comparison of anxiety score of medical personnel between 2 groups of sleep quality.

Figure 1. Anxiety comparison based on Sleep Quality



In figure 1, the anxiety score level is classified as normal, moderate and extreme based on clinical anxiety range in the SAS instrument. Based on the figure, respondents who are at moderate anxiety level are dominated by medical personnel with bad sleep quality (blue line). Meanwhile, almost all medical personnel who have good sleep quality are in the normal range of anxiety level. Table 3 shows the result of comparison analysis based on paired t-test analysis.

Comparison pair	Levene's Test for Equality of Variances		T-test for Equity of Means	
	F	Sig.	Sig. (2-tailed)	Mean Difference
anxiety of "good" SQ - Anxiety of "bad" SQ	2,391	0,129	0,020	5,78788

Comparison test between 2 groups of sleep quality shows a significant difference (p-value/sig. 0.20) with a mean difference of 5,78. Based on the maximum score of anxiety, it can be concluded that HCW with good sleep quality has a 7.35 % lower level of anxiety compared to those who had worse sleep quality..

DISCUSSION

During the pandemic the workload of medical taff in emergency unit had increased(1). They have to deal with the unusual condition which extra effort and energy needed. These condition will potentially increase the mental and physycologicalproblem among medical taff such as anxiety. In this study, the average score of anxiety among HCP in Harjono S Hospital is 32.9 which categorizes into normal range(25). But some medical personnel have a score of more than 45 so they are included in the moderate category on the level of anxiety. Based on data obtained, there are 4 HCP with moderate range of anxiety. Based on the follow-up interview, these anxiety usually caused by the fear of transmitting the disease to their families, the case increased, intensive PPE usage and the more absent number of personnel caused by isolation .

Sleep quality has been widely known to be related to many physical states and physiological functions of humans. Insufficient sleep even become a predictor for clinical burnout (22). Of all the psychiatric disorders associated with insomnia, depression and anxiety is the most common(5).Fredriksen(26) provided longitudinal data to show that sleep loss was a significant predictor of increased depressive feeling and self-esteem. Sleep disturbances encompass various potentially overlapping symptoms and disorders including insomnia, hypersomnia, excessive daytime sleepiness, circadian rhythm disturbance, and extrinsic sleep

disorders (related to insufficient sleep and sleep hygiene). Sleep disturbances, anxiety and depression are common problems that lead to neuropsychological impairment. Neuropsychological evidences suggest that both quality and quantity of sleep are vital to the optimal functioning of brain activity in regulating our emotions (16). Sleep disturbance is diagnostic criteria for depression in the Diagnostic and Statistical Manual of Mental Disorders. Treatment for sleep disorders has been shown to reduce symptoms of depression and anxiety (27).

The goal of this study is to determine whether improved sleep quality could reduce anxiety among medical staff especially in the emergency unit which assumed had more risk to depression and exhaustion. But to get more representative result, advanced study with more samples and instrument might be needed.

The Pittsburgh Sleep Quality Index (PSQI) is a self-rated questionnaire which assesses sleep quality and disturbances over a 1-month time interval. Nineteen individual items generate seven "component" scores: subjective sleep quality, sleep latency, sleep duration, habitual sleep efficiency, sleep disturbances, use of sleeping medication, and daytime dysfunction. The instrument however is only evaluating sleep experiences and not as diagnostic instrument.

In this study, sleep quality is categorized to good and bad instead of numerical scale because it would be easier to determine only "good" or "bad" than increased sleep quality. Though it is suggested that in future research the correlational analysis used within a numeric-numeric scale to obtain more representative results. There is no intervention in this study. Sleep quality is measured by evaluating respondent experience in sleep in the past month. This quasi method may caused the sample size among 2 groups of sleep quality isn't equal. However, the equality number is not required in the independent t-test as long as the number difference is less than 1:4. In the future research, interventions may be needed to obtain an ideal and proportionate sample size for each group.

CONCLUSION

Based on the result concluded that there are significant differences the anxiety score within 2 groups of sleep quality. There are 7.35 % lower scores of anxiety in the workers with good sleep quality compared to those with bad sleep quality.

REFERENCES

1. Shan Y, Shang J, Yan Y, Lu G, Hu D, Ye X. Mental workload of frontline nurses aiding in the COVID-19 pandemic: A latent profile analysis. *J Adv Nurs*. 2021;77(5):2374–85.
2. Stuijzand S, Deforges C, Sandoz V, Sajin C-T, Jaques C, Elmers J, et al. Psychological impact of an epidemic/pandemic on the mental health of healthcare professionals: a rapid review. *BMC Public Health*. 2020;20(1):1–18.
3. Vafaei H, Roozmeh S, Hessami K, Kasraeian M, Asadi N, Faraji A, et al. Obstetrics healthcare providers' mental health and quality of life during COVID-19 pandemic: multicenter study from eight cities in Iran. *Psychol Res Behav Manag*. 2020;13:563.
4. Labrague LJ, de los Santos JA. COVID-19 anxiety among frontline nurses: Predictive role of organisational support, personal resilience and social support. *medRxiv*. 2020;1–25.
5. Pappa S, Ntella V, Giannakas T, Giannakoulis VG, Papoutsis E, Katsaounou P. Prevalence of depression, anxiety, and insomnia among healthcare workers during the COVID-19 pandemic: A systematic review and meta-analysis. *Brain Behav Immun*. 2020;88:901–7.
6. Luo M, Guo L, Yu M, Jiang W, Wang H. The psychological and mental impact of coronavirus disease 2019 (COVID-19) on medical staff and general public--A systematic review and meta-analysis. *Psychiatry Res*. 2020;291:113190.
7. Hu D, Kong Y, Li W, Han Q, Zhang X, Zhu LX, et al. Frontline nurses' burnout, anxiety, depression, and fear statuses and their associated factors during the COVID-19 outbreak in Wuhan, China: A large-scale cross-sectional study. *EclinicalMedicine*. 2020;24:100424.
8. Mak ADP, Wu JCY, Chan Y, Chan FKL, Sung JJY, Lee S. Dyspepsia is strongly associated with major depression and generalised anxiety disorder—a community study. *Aliment Pharmacol & Ther*. 2012;36(8):800–10.
9. Simonetti V, Durante A, Ambrosca R, Arcadi P, Graziano G, Pucciarelli G, et al. Anxiety, sleep disorders and self-efficacy among nurses during COVID-19 pandemic: A large cross-sectional study. *J Clin Nurs*. 2021;30(9–10):1360–71.
10. Nguyen DT, Dedding C, Pham TT, Wright P, Bunders J. Depression, anxiety, and suicidal ideation among Vietnamese secondary school students and proposed solutions: a cross-sectional study. *BMC*

- Public Health. 2013;13(1):1–10.
11. Sunjaya DK, Herawati DMD, Siregar AYM. Depressive, anxiety, and burnout symptoms on health care personnel at a month after COVID-19 outbreak in Indonesia. *BMC Public Health* [Internet]. 2021;21(1):227. Available from: <https://doi.org/10.1186/s12889-021-10299-6>
 12. Zheng R, Zhou Y, Qiu M, Yan Y, Yue J, Yu L, et al. Prevalence and associated factors of depression, anxiety, and stress among Hubei pediatric nurses during COVID-19 pandemic. *Compr Psychiatry*. 2021;104:152217.
 13. Zhan Y, Zhao S, Yuan J, Liu H, Liu Y, Gui L, et al. Prevalence and influencing factors on fatigue of first-line nurses combating with COVID-19 in China: a descriptive cross-sectional study. *Curr Med Sci*. 2020;40(4):625–35.
 14. Dewa CS, Loong D, Bonato S, Trojanowski L. The relationship between physician burnout and quality of healthcare in terms of safety and acceptability: a systematic review. *BMJ Open*. 2017;7(6):e015141.
 15. Jalili M, Niroomand M, Hadavand F, Zeinali K, Fotouhi A. Burnout among healthcare professionals during COVID-19 pandemic: a cross-sectional study. *Int Arch Occup Environ Health*. 2021;1–8.
 16. Urponen H, Partinen M, Vuori I, Hasan J. Sleep quality and health: Description of the sleep quality index. In: *Sleep and health risk*. Springer; 1991. p. 555–8.
 17. Riemann D, Voderholzer U. Primary insomnia: a risk factor to develop depression? *J Affect Disord*. 2003;76(1–3):255–9.
 18. Huang Y, Mai W, Hu Y, Wu Y, Song Y, Qiu R, et al. Poor sleep quality, stress status, and sympathetic nervous system activation in nondipping hypertension. *Blood Press Monit*. 2011;16(3):117–23.
 19. Lavidor M, Weller A, Babkoff H. How sleep is related to fatigue. *Br J Health Psychol*. 2003;8(1):95–105.
 20. Ekstedt M, Söderström M, Åkerstedt T, Nilsson J, Søndergaard H-P, Aleksander P. Disturbed sleep and fatigue in occupational burnout. *Scand J Work Environ & Heal*. 2006;121–31.
 21. Arifah DA, Andarini YD, Dianita R. Occupational Fatigue Based on Work Shift Among Medical Workers at Harjono S Hospital. *J Ilmu Kesehat Masy*. 2019;10(3):199–206.
 22. Söderström M, Jeding K, Ekstedt M, Perski A, Åkerstedt T. Insufficient sleep predicts clinical burnout. *J Occup Health Psychol*. 2012;17(2):175.
 23. Giorgi F, Mattei A, Notarnicola I, Petrucci C, Lancia L. Can sleep quality and burnout affect the job performance of shift-work nurses? A hospital cross-sectional study. *J Adv Nurs*. 2018;74(3):698–708.
 24. Buysse DJ, Reynolds III CF, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28(2):193–213.
 25. Zung WWK. Self-rating anxiety scale. *Arch Gen Psychiatry*. 1971;26:112–8.
 26. Fredriksen K, Rhodes J, Reddy R, Way N. Sleepless in Chicago: tracking the effects of adolescent sleep loss during the middle school years. *Child Dev*. 2004;75(1):84–95.
 27. Freeman D, Sheaves B, Goodwin GM, Yu L-M, Nickless A, Harrison PJ, et al. The effects of improving sleep on mental health (OASIS): a randomised controlled trial with mediation analysis. *The Lancet Psychiatry*. 2017;4(10):749–58.