

MALAYSIAN JOURNAL OF
Medicine and Health Sciences
Vol. 19 No. 4, July 2023

MALAYSIAN JOURNAL OF

Medicine and Health Sciences

Vol. 19 No. 4 July 2022

Malaysian Journal of Medicine and Health Sciences Vol. 19 No. 4, July 2023



Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://medic.upm.edu.my/>

UPM Press
Universiti Putra Malaysia
43400 UPM Serdang
Selangor Darul Ehsan
Malaysia
<http://penerbit.upm.edu.my>



A scientific journal published by Universiti Putra Malaysia Press

About the Journal

The Malaysian Journal of Medicine and Health Sciences (MJMHS) is published by the Faculty of Medicine and Health Sciences, Universiti Putra Malaysia. The main aim of the MJMHS is to be a premier journal on all aspects of medicine and health sciences in Malaysia and internationally. The focus of the MJMHS will be on results of original scientific research and development, emerging issues and policy analyses pertaining to medical, biomedical and clinical sciences. The Malaysian Journal of Medicine and Health Sciences is now indexed in the following data bases: Scopus, EBSCOhost, ISC, and Rubriq.

Editorial Board

Editor in Chief:

Prof. Dr. Normala Ibrahim

Members:

Prof. Dr. Rukman Awang Hamat

Prof. Dr. Christopher Lim Thiam Seong

Prof. Dr. Ching Siew Mooi

Prof. Dr. Sharmili Vidyadaran

Prof. Dr. Loh Su Peng

Assoc. Prof. Dr. Rajesh Ramasamy

Assoc. Prof. Dr. Sethu Thakachy Subha

Assoc. Prof. Dr. Nor Afiah Mohd Zulkefli

Assoc. Prof. Dr. Wan Aliaa Wan Sulaiman

Assoc. Prof. Dr. Ling King Hwa

Assoc. Prof. Dr. Habibah Abdul Hamid

Assoc. Prof. Dr. Zubaidah Nor Hanipah

Assoc. Prof. Dr. Intan Hakimah Ismail

Assoc. Prof. Dr. Ho Yu Bin

Assoc. Prof. Dr. Karmegam Karuppiah

Assoc. Prof. Dr. Omar Ahmed Fahmy Ahmed

Dr. Navin Kumar A/L Devaraj

Dr. Raymond Yeak Dieu Kiat

Dr. Razif Abas@Buang

Dr. Sandra Maniam

Technical Coordinator:

Dr. Tan Kit Aun

Dr. Manraj Singh Cheema

International Advisory Board

Prof. Dr. Cecilia A. Essau

Prof. Dr. David Brendan Price

Prof. Dr. S Suresh Kumar

Assoc. Prof. Dr. Sean Semple

Dr. Martin Wong Chi Sang

Malaysian Journal of Medicine and Health Sciences

Vol. 19(4), July 2023

Contents

EDITORIAL

- Enhancing Occupational Safety and Health for Traffic Police Officers: A Call for Action 1
Karmegam Karuppiah

ORIGINAL ARTICLES

- Malay Version of Asthma Knowledge Questionnaire: A Validation and Reliability Study 3
N. Fafwati Faridatul Akmar Mohamad, Azizah Musa, Faridah Derani, Nor Khamizah Mesiran, Asiah Kassim, Farizah Mohd Hairi, Hafizah Zainuddin
- Risk Perception and Emergency Preparedness Against Flood Affected Participants From the Primary Health Care Centre, Malaysia: A Comparison Between Genders 8
Siew Mooi Ching, Lee Kai Wei, Mari Kannan Maharajan, Hani Salim, Ng Jun Ying, Ng Kar Yean, Aneesa Abdul Rashid, Dhashani A/P Sivaratnam, Nurainul Hana Shamsudin, Imran Ahmad, Vasudevan Ramachandran
- Relationship of Prior Knowledge and Scenario Quality With the Effectiveness of Problem-based Learning Discussion among Medical Students of Universitas Malikussaleh, Aceh, Indonesia 15
Mulyati Sri Rahayu, Sri Wahyun, Yuziani
- Association of Circulating Matrix Metalloproteinase-9 and the Risk of Coronary Heart Disease in Young Smokers 21
Dinesh Nath, Meera Shivasekar, V.M. Vinodhini
- Effect of Addition Tempeh Flour on Calcium, Protein, Water Content and Acceptance in Catfish Meatballs 28
Widyasmara Nur Tammamia, Farida Wahyu Ningtyias, Sulistiyani
- Awareness and Preparedness of Fomite Transmission Among Malaysian Online Shoppers: A Cross-Sectional Survey 34
Hisyam Abdul Hamid, Nor Aslily Sarkam, Mahmathi Karuppanan, Abdul Rafeq Saleman, Noreen Husain
- Effectiveness of Education Using Comic Media on Knowledge About Covid-19 Among Elementary School Students 40
Arif Rohman Mansur, Ira Mulya Sari, Yelly Herien, Meri Neherta, Mei Chan Chong
- Factors Associated with Work-related Musculoskeletal Disorders Risk among Bamboo Craftsman in Mojorejo Village, Ponorogo Indonesia 46
Ratih Andhika Akbar Rahma, Dian Afif Arifah, Eka Rosanti
- Contribution of Snacks and Physical Activities to Elevated Blood Pressure of Adolescent Girls in West Java, Indonesia 53
Mohammad Zen Rahfiludin, Alfi Fairuz Asna, Suroto, Lilik Hidayanti
- Perspective of Primigravida Mother's Experience in Their Birthing Process Utilising HypnoBirthing 61
Thavamalar Paramasivam, Umapathi Mariappan, Chong Mei Chan
- Education Perception, Experience, and Anxiety Level Among Undergraduate Dental Students in Malaysia During the Covid 19 Pandemic 68
Nur Hafizah Kamar Affendi, Nor Faezah Md Bohari, Ameer Razek Bin Salihan, Nur Aishah Binti Rosman
- Recommendation for Improvement in COVID-19 Vaccine Program in Malaysia: An Online Nationwide Qualitative Survey 76
Hana Maizuliana Solehan, Mohd Dzulkhairi Mohd Rani, Nurul Azmawati Mohamed, Muslimah Ithnin, Abdul Rasyid Ariffien, Che Ilina Che Isahak
- Factors Associated with Presence of Human Papillomavirus Infection among Women: Findings from New Cervical Cancer Screening in Kelantan, Malaysia 84
Nur Dalilah Mohd Zin, Tengku Alina Tengku Ismail, Noran Hashim
- Critical Care Nurses's Knowledge and Attitude Towards Sleep Promoting Interventions Among Critically Ill Patients in Intensive Care Unit 93
Maisarah Razali, Rosuzeita Fauzi, Shaharudin Shah Che Hamzah
- Entrepreneurial Intention of Medical Students in Malaysia: Does Socio-Demographic Profile Matter? 103
Kavitha Ashok Kumar, S.M. Ferdous Azam, Ashok Kumar Jeppu3
- Budget Impact Analysis of Ferric Derisomaltose for the Treatment of Iron-Deficiency in Malaysia 110
Huey Yi Chong, June Wai Yee Choon, Wei Chern Ang, Kenneth Kwing Chin Lee

Determinants of Premarital Sexual Behavior Among the Late Adolescents in Pahang, Malaysia <i>Halimatus Sakdiah Minhat, Raja Nurul Najwa Raja Ismail, Lim Poh Ying</i>	117
Thromboelastography Parameter and Its Association with Survival of COVID-19 Patients: A Retrospective Cross-Sectional Study <i>Nita Wiyono, Yetti Hernaningsih, Arifoel Hajat, Paulus Budiono Notopuro, Narazah Mohd Yusoff, Emmanuel Jairaj Moses</i>	124
Risk Factors for Major Adverse Cardiac Events Outcomes in Post Percutaneous Coronary Intervention during Index Admission <i>Muaath Ahmed Hasan Mohammed, Zulkefli Sanip, Zurkurnai Yusof, W. Yus Haniff W. Isa</i>	130
Associations between Sociodemographic, Body Mass Index, Body Image Perception, Physical Activity and Sleep Quality with Mental Health among University Students in Universiti Putra Malaysia during the COVID-19 Pandemic <i>Noraida Omar, Farrah Haliny Kudin</i>	139
Assessment Of Gingival Phenotype At The Aesthetic Zone In A Malay Population Using Probe Test <i>Muhammad Azat Awang@Semail, Abdul Shukur Mustafa, Munirah Yaacob, Juzaily Husain, Noorhazayti Ab. Halim</i>	148
Correlation of the Risk Factors in Degenerative Lumbar Spondylolisthesis with MRI Imaging <i>Paulus Rahardjo, Harry Wahyudhy Utama, Rosy Setiawati, Damayanti Tinduh</i>	153
Subchronic Administration of High-dose Monosodium Glutamate Causes Spatial Memory Dysfunction and Structural Deficits of Rat Hippocampus <i>Muchtar Hanafi, Faizal Muhammad, Amilia Ramadhani, Nurlaela, Zaenal Muttaqien Sofro, Ginus Partadiredja</i>	159
Effect of <i>Moringa oleifera</i> Leaf Extracts on Depression in Rheumatoid Arthritis Patients <i>Afiyah Syifa'ul Ummah, Faizal Muhammad, Yulie Erida Nur Rahmawati, Isa Ridwan, Arief Nurudhin, Eillien Ramadhani Fauzi, Ratih Tri Kusuma Dewi, Yulyani Werdiningsih</i>	166
Cellular Uptake of <i>Catharanthus roseus</i> -Silver Nanoparticles in Human Hepatocellular Carcinoma HepG2 Cells <i>Nur Asna Azhar, Siti Aishah Abu Bakar, Siti Hawa Ngalim, Nor Hazwani Ahmad</i>	171
Preschool Children Feeding Patterns And Early Childhood Oral Health Impact In Tanah Merah, Kelantan, Malaysia <i>Nizamuddin Pardan, Ruhaya Hasan, Norsamsu Arni Samsudin, Wan Muhamad Amir W Ahmad</i>	178
DIMAS (Diabetes Management Support) Application: Mobile-Phone Based Intervention to Support Diabetes Self-Management Activities <i>Annisa Wuri Kartika, Widyatuti Widyatuti, Ety Rekawati</i>	186
Prevalence and Predictors of Anxiety and Depressive Symptoms among Adult Patients with Atopic Dermatitis in Malaysia Tertiary Referral Hospital <i>Nur Hafidah Ishak, Chong Seng Choi, Normala Ibrahim, Firdaus Mukhtar, Umi Adzlin Silim</i>	193
Evaluation of Dental Students' Medical History Records on Hypertension and Diabetes Mellitus at The National University of Malaysia <i>Jun Ai Chong, Fara Azwin Adam, Ang Yee, Laila Azwa Hassan, Hetal Ashvin Kumar Mavani, Rama Krsna Rajandram</i>	201
Attitude Towards Anti-smoking Measures and Its Associated Factors Among Adults in Sub-urban Area, in Kuala Terengganu, Terengganu, Malaysia <i>San San Oo, Myat Moe Thwe Aung, Mohd Salami Ibrahim, Nyi Nyi Naing, Intan Suhana Munira Mat Azmi, Aniza Abd Aziz, San Thitsa Aung</i>	207
Completers Versus Dropouts of A Parent Training Programme and Child Behaviour Improvement In Malaysia <i>Ruziana Masiran, Normala Ibrahim, Hamidin Awang, Lim Poh Ying</i>	214
Public Health Bachelor's Perception of Skills Needed to Work in the Health Office: A Qualitative Study among Alumni from Universitas Negeri Malang <i>Dian Mawarni, Anindya Hapsari, Tika Dwi Tama, Windi Chusniah Rachmawati, Shofiyatul Masyiyah, Nurul Jannatul Firdausi</i>	225
Psychometric Properties of The Indonesian Version of Hypertension Self Care Profile <i>Salami Salami, Soon Siew Choo, Faridah Mohd Said, Angga Wilandika</i>	231
Predictors of Adherence to Personal Preventive Behaviors Among Nursing Students Based on Health Belief Model: Cross Sectional Study During the Second Wave of COVID-19 Pandemic in Indonesia <i>Akbar Satria Fitriawan, Wiwit Ananda Wahyu Setyaningsih, Erni Samutri, Dedi Kurniawan, Fitrio Deviantony, Gatot Suparmanto, Bayu Fandhi Achmad, Eriyono Budi Wijoyo</i>	237
Factors Associated with Diet Quality during COVID-19 Pandemic among Undergraduate Students in Universiti Putra Malaysia <i>Noraida Omar, Dalina Puteri Aripin</i>	247

Morphology of Mandibular Condyle in The Population of Sarawak: A Retrospective Cross-sectional Study Using Digital Panoramic Radiograph <i>Tan Yy Jean, Lim Woei Tatt, Lee Sie Wei, Shim Chen Kiong, Mohamad Adam Bujang</i>	258
Fear of Covid-19 and Burnout Among Healthcare Providers in Malaysia: Is Resilience a Missing Link? <i>Siew-Mooi Ching, Ramayah Thurasamy, Ai Theng Cheong, Anne Yee,, Poh Ying Ling, Irmis Ismail Zarina, Kai Wei Lee, Jun Ying Ng, Roſina Abdul Rahim, Mohd Khairi Mohd Noor, Chang Li Cheng, Ahmad Iqmer Nashriq Mohd Nazan, Hafizah Md Salleh, Noor Hasliza Hassan</i>	265
Dietary Inflammatory and Its Association with Cognitive Frailty Among Community-Dwelling Older Adults in Klang Valley <i>Felicia Bong Jun Fae, Rasyidah Abdul Rahman, Nurul Hidayah Md Fadzil, Suzana Shahar</i>	273
The Level of N-Carboxymethyllysine and C-Reactive Protein in Type 2 Diabetes Mellitus and it's Association with HbA1c in Diabetic Nephropathy <i>Mohd Zulhafiz Mat Rahim, Wan Norlina Wan Azman, Najib Majdi Yaacob, Wan Mohd Izani Wan Mohamed, Noorazliyana Shafii</i>	282
Nurses' Perception of Interprofessional Collaboration in Hospitals in the Maldives <i>Aishath Selna, Zulhabri Othman, Jacqueline Tham, Adam Khaleel Yoosuf</i>	290
SYSTEMATIC REVIEW	
Dichoptic Vision Therapy in Adults with Anisometropic Amblyopia: A Systematic Review <i>Rituparna Ghoshal, Dipanwita Ghosh, Somnath Ghosh</i>	297
Effectiveness of Lumbar Stabilization Exercise on Pain, Disability and Functional Performance among Older People with Low Back Pain: A Systematic Review <i>Zarina Zahari, Siti Aishah Shaiful Azan, Naim Faiz Ahmad Imran</i>	307
Improving Foot Self-care Practices Through Health Education Intervention Programs Among Diabetic Patients: A Systematic Review <i>Sampson Emilia Oluchi, Rosliza Abdul Manaf, Suriani Ismail, Hayati Kadir Shahar, Theophilus Kachidelu Udeani</i>	315
REVIEW ARTICLES	
The Role of Oxidative Stress and Inflammation in Prediabetes: A Review <i>Fatma S.A Saghir, Farrah Shafeera Ibrahim, Zulkhairi Amom, Lekhsan Othman</i>	326
Application of Sendai Framework for Disaster Risk Reduction in Food Security During Flood Disaster in Malaysia: A Narrative Review <i>Muhammad Ikhwan Mud Shukri, Nazmi Ainaa Azmi, Chandramalar Kanthavelu, Nur Za'Imah Zainal, Rosliza Abdul Manaf, Suhainizam Muhamad Saliluddin</i>	332
Effect of Paint Exposure Among Paint Workers and DNA Damage: A Scoping Review <i>Afiqah Saironi, Normah Awang, Anuar Ithnin, Nurul Farahana Kamaluddin, Farah Wahida Ibrahim, Kok Meng Chan</i>	342
Barriers and Facilitators Factors to Uptake of Cervical Cancer Screening Among Women in Low- and Middle-income Countries: A Narrative Review <i>Sarah Ahmed Al-Oseely, Rosliza Abdul Manaf and Suriani Ismail</i>	354
LETTER TO THE EDITOR	
Embracing Big Data in Medical Education in Malaysia: Overcoming Challenges and Seizing Opportunities <i>Syahrul Nizam Junaini, Keng Sheng Chew</i>	360
COMMENTARY	
Augmented & Virtual Reality via High Mounted Display: Commentary to Cybersickness Questionnaires <i>Nur Izah Ab Razak, Razif Abas, Rahmita Wirza O.K. Rahmat</i>	361
CASE REPORT	
Reviewing the Compounding Risks for Tuberculosis Drug-Induced Liver Injury in a Patient with Transfusion Dependent Thalassaemia <i>Nursafinas Rofii, Ummi Nadira Daut, Su Yin Lau</i>	365
Prostatic Metastasis Mimicking a Ligamentum Flavum Cyst in Thoracic Spine: A Case Report <i>Chee Leong Choong, Chee Kean Lee, Kularaja Thamasegaran, Ker Woon Choy</i>	368

ORIGINAL ARTICLE

Factors Associated with Work-related Musculoskeletal Disorders Risk among Bamboo Craftsman in Mojorejo Village, Ponorogo Indonesia

Ratih Andhika Akbar Rahma, Dian Afif Arifah, Eka Rosanti

Program Studi Keselamatan dan Kesehatan Kerja, Fakultas Ilmu Kesehatan, Universitas Darussalam Gontor, Jl. Raya Siman Km 5-6, Demangan, Siman, Kab. Ponorogo, Jawa Timur, Indonesia

ABSTRACT

Introduction: Work-related Musculoskeletal disorders (WMSDs) becomes the one of most-occupational problem and highly related to the individual and work factors. Eighty percent (80%) workers in the woven bamboo industry in Mojorejo Village dominated by middle aged workers (>40 years) that most of their producing activities are manually handled, without using machine or modern technology. Incompatibility work station, long working hours and repeated movement aggravated the WMSDs risk. This study aims to describe the WMSDs risk and determine its factors based on individual characteristics. **Methods:** There are 58 workers as respondents in this cross-sectional study. Data were collected and analyzed using quantitative method. Posture Exposure (PE) were measured using the QEC instrument. While the WMSDs risk were measured using PLIBEL. Individual factors (PE, Age, Working Period) related to WMSDs were analyzed using Multinomial Logistic Regression (95% CI). The Chi-square test were also used to compare PE rates among WMSDs risk levels. **Results:** Most respondents (48.3%) with 51-70% PE rate have a high level of WMSDs risk. While 27 % respondents with PE rate >70% have a very high level. The neck is the region most exposed to the WMSDs (score 15.5/very high level risk). PE rate is the only factor that significantly related to WMSDs risk (sig. = 0.043; Rsquare = 0.529; r =0.703). While the age and working period has no significant correlation (Sig = 0.859 and 0.851 respectively). **Conclusion:** Based on the result, immediate improvement required to be implemented. It is suggested to redesigned work stations adjusted to the workers posture and position.

Malaysian Journal of Medicine and Health Sciences (2023) 19(4):46-52. doi:10.47836/mjmhs19.4.9

Keywords: WMSDs, Musculoskeletal Disorders, Work Posture, Bamboo Craftsmen, Woven Bamboo Industry

Corresponding Author:

Dian Afif Arifah, M.Kes

Email: dianafif@unida.gontor.ac.id

Tel: +62 856 48185169

INTRODUCTION

Musculoskeletal Disorders refer to symptoms caused by disorders or damage to the joints, ligaments or muscles. There are various types of MSDs depends on the region affected or injured. Over-stretching, heavy loads and repetitive activities are the most common causes for most types of MSDs (1). According to Graveling (2), occupational activity recognized as the major factor contributed to MSDs risk due to the intense or long-duration exposure in a human daylife activity.

Work-related Musculoskeletal disorders (WMSDs) is a term used to describe MSDs caused by work activities. WMSDs accumulatively have a major impact on the company and the worker's productivity (1). It is known to be the second leading cause of short-term or temporary

illness after the common cold in the workplace (2). Based on Summers (3), WMSDs contribute to absenteeism in many working populations and becoming the one of most-occupational problem due to increased compensation and health costs, reduced productivity, and lower quality of life of the workers.

Work posture known to be most related to the risk of WMSDs (4). Work posture is an accumulation of position, weight of the load lifted, manual handling, repeated movement and static or dynamic position of the workers. Poor position, over-stretched muscles, heavy, repetitive, and prolonged lifting can exceed the tolerable tissue stress limit, causing injury due to the overactivity or imbalance (5). Kristensen's research (6) shows that there is an evidences that changing position frequently could reducing a risk of owork related musculoskeletal problems. Static position have been significantly to be much more taxing than dynamic forces because of holding the body in the same position. During a static position, more than 50% of the body's muscles contract to sustain the position against gravity (7-9).

However, most musculoskeletal disorders (MSDs) are characterized as multifactorial (1). Based on Da Costa (5), there are biomechanical, psychosocial and individual risk factors associated to WMSDs. Biomechanical factors including e heavy lifting, manual handling, repetitive movement, and every kind of awkward static or dynamic postures. Psychosocial fators identified as stress, low motivation, and high work dissatisfaction. While the individual risk factors are identified by older age, female gender, longer working period, smoking, obesity, and comorbidities. Every working activity usually has specific working posture that affect specific body region to WMSDs risks. In order to minimizing WMSDs effect, several programs and policy continuously improve in the most large companies. Meanwhile workers in Small to Medium Enterprises (SMEs) known tobe less prioritized as implementation subject in reducing WMDs risks (10).

One of the SMEs with the high risk factors of WMSDs is traditional creative industries due to the most of its productions are manually handled. Culture-based industries as part of the creative industries has a significant contribution to national economic growth in Indonesia (11). According to the data from the Ministry of Tourism and Creative Economy, in 2013, the contribution of the creative industry to GDP reached Rp 578 trillion where 11.8 million people were involved in creative industry employment and 3.1 million people of them hired in the craft industry (12). In some creative industry, not all products are mass-produced. Some of them are hand-crafting with a high artistic feel in the home industry. In Indonesia, home industry is mostly carried out by people in rural areas, where they only have small capital, and also do not have much network in their business (13).

One of the popular creative industry in Indonesia is woven bamboo-based handicrafts. The craft of woven bamboo is very familiar, frequently used and continuously produced because traditional packaging are more interested recently. The bamboo material is cheap and could be simply planted in all over rural areas (14). Producing bamboo handcraft has its own complexity since every part must be carefully made, mostly using manual equipment (15). Due to the less-applied technology, the operational production usually takes longer process depending on worker’s skill and expertise.

In the Mojorejo village Jetis Ponorogo, East Java, woven bamboo industry has succeeded to employ local workers and improve the regional economy within 10 years. There are 40 families who were actively participated in the production process of woven bamboo crafts in Mojorejo Village. The products produced are besek, steamer, capil and chicken cage etc (16). Eighty percent (80%) workers in this industry dominated by middle aged workers (>40years old). Based on Diannita (16) working as a woven craftsman is less attractive to young people

because their motivation is low to produce traditional products. The workers mostly work in a sitting position and make handicraft products for approximately 10 hours a day using makeshift tools and facilities. The 6 main variety of bamboo crafts produced in the industry including : Besek, cone steamer, basket, farmers hat and chicken cage. No job divission or activity so that the workers randomly did the entire stage to make a woven bamboo product. After chopped bamboo tree were obtained, the production steps described in the table 1.

Table 1: The Production Process

Production Steps	Activity	Duration
Making sharpened and mashed bamboo sheets	Slice bamboo into sheets, mashed and sharpened it with a knife and sandpaer based on the size required in the products	± 2 hours/ 500 sheets
Making hard sheet product frame	Slice bamboo into hard sheets as a product frame for basket, hat and chicken cage	± 2 hours/ 20 sheets
Weaving bamboo sheets	Manually waving bamboo sheets into various form of the product	± 5 hours/ 20 woven sheets
Tie up woven sheet with frame	Manually tie woven sheet with a frame using ropes	± 10 minutes/ product

In a day, an average worker can make 10-15 woven bamboo products, depending on the type and complexity of the product produced. The salary obtained based on the amount and type of product they can produce. There are no binding regulations, facilities or safety guarantees because the workers are voluntarily participated (17).

Based on biomechanical factors, statically bending in the sitting position and frequently involved manual handling are the common activity in the traditional woven bamboo craft industry. Working position and activities in the woven bamboo craft industry ilustrated in the Figure 1. Work that frequently supported by hand activity increase risk carpal tunnel syndrome, shoulder and neck pain (17, 18). While static sitting and bending position are biomechanical factors causing shoulder, neck and low back pain (19).

Based on the individual factors, working in the older age and long working period are two common characteristics among respondents. Age-related musculoskeletal disorders (MSDs) are characterized by loss of muscle strength, bone fragility, loss of cartilage strength, and reduced elasticity of ligaments (20). While working period related to the repetitive activities that continuously carried out for a long time. Static workloads may be acceptable in normal working hours (8 hours), but workers with 10 working hours per day, even with normal working week often proved to be an important factor regarded the developing of WMSD (21).

This study aims to describe the risk of WMSDs and determine its factors represented by working posture, age



Figure 1. The activity in the woven bamboo industry: (1) Making hard-shaped product frame, (2) Making sharpened and Mashed-bamboo sheets, (3) Waving bamboo sheets

Source : Muslih's documentation (18)

and working period among woven bamboo craftsmen in Mojorejo Village. The results of this study could represent how workers in traditional creative industries are exposed to occupational risks that may affect their quality of life. It is hoped that further research or programs could determine the appropriate interventions to be applied based on the problems faced by SME's workers.

MATERIALS AND METHODS

Data collection and Respondents

The observational study with cross-sectional method was conducted to obtain entire data in this research quantitatively. The study involved all of woven bamboo craftsmen in Mojorejo Village who were joined in the Industry as a population consisting of 67 workers. Selected respondent qualify as a sample should been working for at least 5 years and had no injuries or chronic diseases in the last 10 years.

Even though targeting entire population, the missing in data collection was anticipated with require minimum sample size following the formula for finite population :

$$n = \frac{N}{1+(Ne^2)} , (22)$$

with (n = minimum sample size required; N = total population; and e = tolerable error rate = 5% = 0,05).

so that the minimum sample size estimated as :

$$n = \frac{67}{1 (67 \times 0.05^2)} = 57.38 \sim 58 \text{ respondents}$$

Due to the missing in data collection, there are 58 workers who were selected as respondents in this study. They were informed about the objectives and procedures of the study and voluntary participated as respondent by signing an informed consent.

Ethical Issue

This research has complied with the ethical guidelines set by the Ethical Research Publishing Committee and has been approved by the Review Board of the Harjono S Hospital Ethics Committee with the registered number 0054213502121242022100400/X/KEPK/2022. All respondents are voluntarily participate after they get a full explanation about procedures, benefits and consequences of the research so that there is no ethical issues in this study.

Variables and Instruments

The Posture rating score were measured using the Quick Exposure Check (QEC) questionnaire. Risk rating score obtained based on the complaints experienced in eight (8) components (Back, Shoulder/arm, Wrist/hand, Neck, Stable base, Vibration, Work pace and Environment) which is categorized into Low, Moderate, High and Very High risk. Rating score for each category described in the table II.

Table II: Risk Rating levels each region and activity in QEC

Exposure	Risk Rating (RR)			
	Low	Moderate	High	Very High
Body region				
Back	10-20	21-30	31-40	41-56
Shoulder/Arm	10-20	21-30	31-40	41-56
Wrist/Hand	10-20	21-30	31-40	41-56
Neck	2-6	7-10	11-14	15-18
Activity				
Stable Base	1	4	9	
Vibration	1	4	9	-
Work Pace	1	4	9	-
Environment	1	4	9	16

While the Work Related Musculoskeletal Disorders (WMSDs) risk were measured using PLIBEL instrument. In the measurement process, the work condition and body region exposed (neck shoulder, upper back, elbows, forearm, hands, feet, knees and hips, dan low back) being evaluated to gain exposure risk in the percentage rates then categorized into four (4) level of exposure rates (≤40; >40 - 50 ; >50 - 70 ; and >70).

Validity and Reliability Test

There are no validity and reliability test for QEC since it objectively assumed and measured by observator team. The item validity of the PLIBEL was tested involving a

separate group consisting 30 samples with the similar characteristic as respondents in the study. Item validity determined based on the Pearson correlation between item to the total answers then compared to the r-table at $df = 28 (n-2)$ and 95% CI. There were no dropped out item (pearson correlation score was more than (\geq) r-table) so that all items are qualify as valid.

The instrument reliability test also conducted to measure answer consistency among respondents based on the Cronbach Alpha score. In this study, the Cronbach Alpha score obtained is 0.868 or > 0.7 (23) so that the instrument interpreted as sufficiently reliable.

Data Analysis

Data were both descriptively and analytically processed. Microsoft Excel and Statistical Package for Social Sciences Software (SPSS) version 24.0 was used in most steps of data analysis. The multivariate analysis to determine factors (PE, Age and Working Period) related to WMSDs risk were analyzed using Multinomial Logistic Regression Model with 95% of Confidence Interval (CI). The Chi-square test also used as non-parametric test to compared PE rate among WMSDs risk levels.

RESULTS

Respondent’s individual characteristics were described by age and work period in years. Table III describe the respondents distribution based on age and work period.

Respondent’s age ranged from 35 to 76 years and the average respondents were 59.1 years. Based on the work period, the respondents has been working for at least 5 years and 8.6 years in average. Table IV showed the QEC measurement result described average score and level of the MSDs risk based on exposure source among 58 workers. The result of the table IV showed that each body region exposed to the risk of MSDs at least in the moderate level. The neck is the region most exposed to the risk of MSDs. Besides, table IV also shows descriptively the Posture Exposure rate and WMSDs risk level each category. Action level described the action needed based on the risk level. Posture Exposure rate shows a percentage of body regions are exposed to un-ergonomic position.

Table III: Descriptive data of the Age and Work Period

	Variables	
	Age (years)	Work Period (years)
Average	59.10	8.81
Min	35.00	5.00
Max	76.00	17.00
S.Deviation	9.26	2.97

Table IV: Muskuloskeletal Disorders Risk based on Body Posture, Activity and Risk Level of Exposure

Region and Activity Exposed	Score	Risk Level	
Body Region			
Back	32.2	High	
Shoulder/ Arm	30.8	Moderate	
Wrist/ Hand	23.5	Moderate	
Neck	15.5	Very High	
Activity			
Stable Base	1.0	Low	
Vibration	1.0	Low	
Work Pace	3.7	Low	
Environment	2.5	Low	
Level/ Rate	Action Level	n	%
Posture exposure (%)			
≤ 40	Acceptable	0	0.0
$> 40 - 50$	Investigate further	0	0.0
$> 50 - 70$	Investigate further and change soon	33	56.9
>70	Investigate and change immediately	25	43.1
MSDs Risks			
Low	No action needed	0	0.0
Moderate	Action might be needed	5	8.6
High	Action needed immediately	37	63.8
Very High	Action needed urgently	16	27.6

In the posture exposure rate, it is known that the most of the respondents (56.9%) have 3rd level (51-70% exposure rate) of posture exposure and the rest of it (43.1%) have a higher level. The table shows that all of the workers have at least 51% or higher posture exposure rate of non-ergonomic position so that further investigation needed to determine change. While based on the MSDs Risks measured in PLIBEL, most workers have a high level (63.8%) of subjective MSDs risk level. There are 27.6% of respondents who have a very high level of MSDs risk level so that the actions are urgently needed.

Table V describe WMSDs risk factors based on the Posture Exposure, Age and Working Period. There are 3 predictor variables partially correlated to WMSDs in categorical score (1-4) including Posture Exposure, Age and Working period. Based on significant value, it is known that among all factors correlated to WMSDs, only Posture Exposure (Sig./p-value < 0.05) that is significantly qualify as a model predictor for WMSDs

Table V: Logistic Regression Model based on predictors to WMSDs

Variable	Pseude R-square	Sig. (2-tailed)	Model fit Final (Sig. 2-tailed)
(constant)	0.529	0.00*	0.00
Posture Exposure		0.043*	
Age		0.859	
Working Period		0.851	

* $\alpha < 0.05$ ** $\alpha < 0.01$

score. Meanwhile the Age and Working period has no significant effect to WMSDs risk (Sig > 0.05). Pseudo R square interpreted as the contribution of Posture Exposure to Increase WMSDs risk is 0.529 or 52.9 %. Since the posture exposure and WMSDs risk are often scaled as categorical, further analysis is non-parametrically describe the crosstabulation between PE and WMSDs.

It is known that the most respondents (48.3%) with 51-70% posture exposure rate have a high level of WMSDs risks. While 27 % respondents with an exposure rate >70% have a very high level of MSDs risks. The table VI shows the pearson chi-square analysis (with the 95 % CI) between two variables. The p-value score of 0.00 (<0.05) shows that there are significant relationship between posture exposure rate and WMSDs risk variables with a correlation score 0.703. Likelihood ratio interpreted as the possibility of an increased WMSDs risk due to an increased posture exposure rate is 38.24 %.

Table VI: The chi-square test result between Posture Exposure and WMSDs risk

Indicators	Value
Pearson Chi-Square (p-value)	0.000
Likelihood Ratio	38.240
Spearman correlation	0.703
N	58

DISCUSSION

The results of this research descriptively indicate that there are non-ergonomic work stations and require immediate improvements. Based on the results of measuring WMSDs risk using the PLIBEL instrument shows that 63.8% workers need immediate action and the rest of 27.6% need urgent action, while the measurement of work posture using the QEC instrument shows that all respondents need an immediate improvement thorough investigation of working conditions. The findings in this study are very important as a management reviews to determine work stations improvements.

The neck and the back is the body region most exposed to MSDs risk. It might correlated to the frequently bending position in e long duration while working on making crafts. Wijayanti's (24) research also shows that handcraft workers experience severe neck complaints due to being in a bent position for too long while

working. Schibye et al (25) studied the lack adjustment of sitting position is a risk factor highly correlated with subjective neck symptoms. Bernard (26) proved some evidence that repetitive hand or arms movements and static postures involving the neck or shoulder muscles contributed to the neck pain and disorders.

The high risk of musculoskeletal disorders among woven bamboo workers is also described in Sriagustini's research (27) which shows that the main ergonomic hazards for workers is non-ergonomic work station and lack of individual adjustment for ergonomic. Nugroho (28) conducted a similar study aimed at reducing WMSDs risk in the rattan weaving process at the Anggun Rattan UKM. One of the results of this research is that the weaving process in Anggun Rattan UKM has a high risk and needs to be improved. The results of the Nordic Body Map also indicate that the operator has musculoskeletal disorders, especially in the neck, shoulders, back and thighs.

The results of observations in this study indicate that all workers work in a static position with a narrow range of motion and perform repetitive movements continuously for approximately 5 hours. According to Carter (29) , if the muscles receive static loads repeatedly for a long period of time, it can cause complaints in the form of damage to joints, ligaments and tendons. These complaints are usually referred to as MSDs or injuries to the musculoskeletal system. Based on the study results which analyzed the posture variable as the only factor affecting WMSDs, it showed that the correlation between posture and WMSDs risk was strong ($r = 0.70$). WHO (x) determine main factors that are directly associated with MSDs among workers including: heavy loads, high force exertion, working in unfavorable body postures, monotonous repetitive movements, long-lasting loading and physical environmental conditions (7).

High muscle strength exertion affect in overstretching of used muscle tissue. High-intensity exertion are active within the body tissues particularly during lifting or carrying heavy objects (30). In addition, pushing, pulling, holding or lifting excess weight causes high-intensity muscle stretching problems. Handling heavy materials for over a long duration caused musculoskeletal failures if performed for several years (31). WMSDs risk in this study may also caused by frequently-repeated movements for a long work duration and long period, even if the weight of the objects handled or the forces produced are low (7). Under such condition, the same muscle region are strained for long periods or with a high frequency. Early fatigue, pain and possible injuries are the consequences.

Muscle load in a static posture is a condition in which muscles are tensed continuously for a long time to hold a certain posture. A characteristic of static muscle loads is that the muscles contract without the appropriate joint

movement being followed. If the muscles do not have the opportunity to rest during the condition, muscle fatigue can occur even at low strength levels. In addition, the static load causes a lack of blood circulation in the muscles (32). Monotonous repetitive movements with or without lifting for long periods of time can cause musculoskeletal disorders. Repetitive work occurs when the same body part is repeatedly activated and there is no possibility of at least a brief period of relaxation, or movement variation is impossible (33). Physical environmental factors such as temperature and climatic conditions can exacerbate the risk of musculoskeletal disorders. Furthermore, psychosocial influences such as unfavorable work demands, facilities, and support from coworkers can add to the effect of physical and psychological tension (34). A risk for disorders of the musculoskeletal system appears if the load and the functional capacity of the worker are not balanced (7). However, MSDs is known to be multifactorial which is difficult to determine overall factors in a single study. It is important to integrate data from different studies on factors that contribute to WMSD. In this way, the risk factors and level of evidence of their association with each WMSD can be evaluated.

This research might be represent condition among woven bamboo craftsmen that needs a further study in another factors, since the various work condition in every working environments. The basic principle of ergonomic is to create an appropriate balance between the requirements of the work and the capacity of the working person, by either adapting the work to the workers by design of the respective work.

CONCLUSION

The result of this study prove that the most of the workers of woven bamboo packaging in Mojorejo Village, Jatis District, Ponorogo Regency experienced musculoskeletal complaints and need an immediate improvements. The neck is the region most exposed to the MSDs risk (score 15.5/very high level risk). Based on the result, work stations need to be redesigned adjusted to the workers posture and position. The finding in this research also proved that there are several occupational problems in the SMEs. Occupational health and safety management system policies should be rearranged so it could cover the entire working community, including workers in the SME

REFERENCES

1. Daneshmandi H, Choobineh AR, Ghaem H, Alhamd M, Fakherpour A. The effect of musculoskeletal problems on fatigue and productivity of office personnel: a cross-sectional study. *J Prev Med Hyg.* 2017;58(3):E252-E258.
2. Yelin EH, Felts WR. A summary of the impact of musculoskeletal conditions in the United States. *Arthritis Rheum.* 1990;33(5):750-755. doi:10.1002/art.1780330520.
3. K. Summers, K. Jinnett, and S. Bevan, "Musculoskeletal disorders, workforce health and productivity in the United States," The center for workforced health and performance. London: Lancaster university, 2015.
4. Arifah DA, Basri AA. Stretching Exercise to Reduce Musculoskeletal Pain among X Bakery ' s Workers. *International Journal of Public Health Science* 2021; 10(3):544-550. doi: 10.11591/ijphs.v10i3.20877
5. da Costa BR, Vieira ER. Risk factors for work-related musculoskeletal disorders: A systematic review of recent longitudinal studies. *Am J Ind Med.* 2010;53(3):285-323. doi:10.1002/ajim.20750
6. Juul-Kristensen B, Hansson GA, Fallentin N, Andersen JH, Ekdahl C. Assessment of work postures and movements using a video-based observation method and direct technical measurements. *Appl Ergon.* 2001;32(5):517-524. doi:10.1016/s0003-6870(01)00017-5
7. WHO and others, "Preventing musculoskeletal disorders in the workplace," Protecting workers 'Health series, no. 5, 2003.
8. Cimelli SN, Curran SA. Influence of turnout on foot posture and its relationship to overuse musculoskeletal injury in professional contemporary dancers: a preliminary investigation. *J Am Podiatr Med Assoc.* 2012;102(1):25-33. doi:10.7547/1020025
9. M. Anghel, V. Argeanu, C. Talpo, and D. Lungeanu, "Musculoskeletal disorders (MSDS) consequences of prolonged static postures," *J Exper Med Surg Res*, vol. 4, pp. 167–172, 2007.
10. V. Y. Siong, J. Azlis-Sani, N. H. M. Nor, M. N. A. M. Yunus, J. A. Boudeville, and S. Ismail, "Ergonomic assessment in small and medium enterprises (SMEs)," *Journal of Physics Conference Series* 2018;1049(1):012065. doi: 10.1088/1742-6596/1049/1/012065
11. F. Z. Fahmi and S. Koster, "Creative industries and regional productivity growth in the developing economy: Evidence from Indonesia," *Growth Change* 2017; 48(4):805–830. doi: 10.1111/grow.12212
12. kemenperin, "Kontribusi PDB Ekonomi Kreatif Ditargetkan 7.5% (Ministry of Industry: Contributor of Bruto Domestic Product to Creative Economy is Targeted by 7.5%)," 2015.
13. S. Budidarsono, A. Susanti, A. Zoomers, and others, "Oil palm plantations in Indonesia: The implications for migration, settlement/resettlement and local economic development," *Biofuels-economy, environment and sustainability*, pp. 173–193, 2013.
14. Y. Sofiana, C. O. Sylvia, and M. Purbasari, Potential of bamboo as material for furniture in

- rural area in Indonesia. *Journal of Computational and Theoretical Nanoscience* 2017; 23(1):263-266. doi: 10.1166/asl.2017.7153
15. Prihatini J. The Challenges of Small Industry of Woven Bamboo Craftsmen to Meet Industrial 4.0 in District of Majalengka, West Java, Indonesia, in ICEASD & ICCOSED 2019: International Conference on Environmental Awareness for Sustainable Development in conjunction with International Conference on Challenge and Opportunities Sustainable Environmental Development, ICEASD & ICCOSED 2019, 1-2 April 2019, 2019, p. 378. doi: 10.4108/eai.1-4-2019.2287242
 16. R. Diannita, M. Muslih, N. Wijayanti, and U. Fatayati, "Participatory Rural Appraisal (PRA) For The Development Of Safe Weaving Bamboo Production Through Community Empowerment To Improve Health And Safety Of Bamboo Weaving Craftsmen In Mojorejo, Ponorogo, East Java, Indonesia," *International Journal Of Community Service (IJCS)*, 2021; 1(3):243–252, 2021. doi: 10.51601/ijcs.v1i3.44
 17. Pourmemari MH, Heliuvaara M, Viikari-Juntura E, Shiri R. Carpal tunnel release: Lifetime prevalence, annual incidence, and risk factors. *Muscle Nerve*. 2018;58(4):497-502. doi:10.1002/mus.26145
 18. Baldasseroni A, Tartaglia R, Carnevale F. Rischio di sindrome del tunnel carpale in alcune attività lavorative (The risk of the carpal tunnel syndrome in some work activities). *Med Lav*. 1995;86(4):341-351.
 19. Hasegawa T, Katsuhira J, Oka H, Fujii T, Matsudaira K. Association of low back load with low back pain during static standing. *PLoS One*. 2018;13(12):e0208877. Published 2018 Dec 18. doi:10.1371/journal.pone.0208877.
 20. Gheno R, Cepparo JM, Rosca CE, Cotten A. Musculoskeletal disorders in the elderly. *J Clin Imaging Sci*. 2012;2:39. doi:10.4103/2156-7514.99151.
 21. Dinar, A., Susilowati, I. H., Azwar, A., Indriyani, K., & Wirawan, M. . Analysis of Ergonomic Risk Factors in Relation to Musculoskeletal Disorder Symptoms in Office Workers. *KnE Life Sciences*, 2018;4(5), 16–29. doi:10.18502/kl.v4i5.2536
 22. Potthoff, Richard F., et al. 'Equivalent Sample Size' and 'Equivalent Degrees of Freedom' Refinements for Inference Using Survey Weights Under Superpopulation Models. *Journal of the American Statistical Association*, 1992;87(418): 383–96. doi:10.2307/2290269.
 23. Nunnally JC, Bernstein IH. *Psychometric theory* (3rd ed.). New York: McGraw-Hill, 1994.
 24. E. W. Wijayati, "Risiko Postur Kerja Terhadap Keluhan Subyektif Nyeri Leher Pada Pekerja Industri Kerajinan Kulit," *JUMANTIK (Jurnal Ilmiah Penelitian Kesehatan)*, 2020; 5(1):56–64. doi: 10.30829/jumantik.v5i1.5891
 25. Schibye B, Skov T, Ekner D, Christiansen JU, Sjøgaard G. Musculoskeletal symptoms among sewing machine operators. *Scand J Work Environ Health*. 1995;21(6):427-434. doi:10.5271/sjweh.58
 26. B. P. Bernard and others, "Musculoskeletal disorders (MSDs) and workplace factors: A critical review of epidemiologic evidence for work-related musculoskeletal disorders of the neck, upper extremity, and low back. US Department of Health and Human Services," National Institute for Occupational Safety and Health (NIOSH), Cincinnati, OH, 1997.
 27. I. Sriagustini and T. Supriyani, "A Analisis Bahaya pada Pengrajin Anyaman Bambu," *Faletehan Health Journal*, vol. 8, no. 03, pp. 223–230, 2021.
 28. A. W. S. NUGROHO, "PERBAIKAN POSTUR KERJA UNTUK MENGURANGI KELUHAN MUSKULOSKELETAL PADA PROSES PENGANYAMAN ROTAN DI UKM ANGGUN ROTAN," *UAJY*, 2019.
 29. Carter JB, Banister EW. Musculoskeletal problems in VDT work: a review. *Ergonomics*. 1994; 37(10):1623-1648. doi: 10.1080/00140139408964941
 30. Gearhart RF Jr, Goss FL, Lagally KM, et al. Ratings of perceived exertion in active muscle during high-intensity and low-intensity resistance exercise. *J Strength Cond Res*. 2002;16(1):87-91.
 31. Hogan DA, Greiner BA, O'Sullivan L. The effect of manual handling training on achieving training transfer, employee's behaviour change and subsequent reduction of work-related musculoskeletal disorders: a systematic review. *Ergonomics*. 2014;57(1):93-107. doi:10.1080/00140139.2013.862307
 32. Jagannath M, Balasubramanian V. Assessment of early onset of driver fatigue using multimodal fatigue measures in a static simulator. *Appl Ergon*. 2014;45(4):1140-1147. doi:10.1016/j.apergo.2014.02.001
 33. Hostens I, Ramon H. Assessment of muscle fatigue in low level monotonous task performance during car driving. *J Electromyogr Kinesiol*. 2005;15(3):266-274. doi:10.1016/j.jelekin.2004.08.002.
 34. Arifah DA, Andarini YD, Dianita R. Occupational Fatigue Based on Work Shift Among Medical Workers at Harjono S Hospital," *Jurnal Ilmu Kesehatan Masyarakat*, 2019; 10(3): 199–206. doi:10.26553/jikm.2019.10.3.199-206